






MOCK KCET Physics QP SET-3

Question 1: A particle shows the distance-time curve as shown in the figure. The maximum instantaneous velocity of the particle is around the point.



1. a. P
2. b. S
3. c. R
4. d. Q

Question 2: Which of the following graphs correctly represents the variation of 'g' on the Earth?



Question 3: A cup of tea cools from 65.5°C to 62.5°C in 1 minute in a room at 22.5°C . How long will it take to cool from 46.5°C to 40.5°C in the same room?

1. a. 4 minutes
 2. b. 2 minutes
 3. c. 1 minute
 4. d. 3 minutes
-

Question 4: The dimensions of the ratio of magnetic flux (ϕ) and permeability (μ) are

1. a. $[\text{M}^0\text{L}^1\text{T}^0\text{A}^1]$
 2. b. $[\text{M}^0\text{L}^{-3}\text{T}^0\text{A}^1]$
 3. c. $[\text{M}^0\text{L}^1\text{T}^1\text{A}^{-1}]$
 4. d. $[\text{M}^0\text{L}^2\text{T}^0\text{A}^1]$
-

Question 5: A mass 'm' on the surface of the Earth is shifted to a target equal to the radius of the Earth. If 'R' is the radius and 'M' is the mass of the Earth, then work done in this process is

1. a. $mgR / 2$
 2. b. mgR
 3. c. $2 mgR$
 4. d. $mgR / 4$
-

Question 6: First overtone frequency of a closed pipe of length ' l_1 ' is equal to the 2nd harmonic frequency of an open pipe of length ' l_2 '. The ratio $l_1 / l_2 =$

1. a. $3 / 4$
 2. b. $4 / 3$
 3. c. $3 / 2$
 4. d. $2 / 3$
-

Question 7: The resistance $R = V / I$ where $V = (100 \pm 5) \text{ V}$ and $I = (10 \pm 0.2) \text{ A}$. The percentage error in R is

1. a. 5.2%
2. b. 4.8%
3. c. 7%
4. d. 3%

Question 8: A block rests on a rough inclined plane making an angle of 30° with the horizontal. The coefficient of static friction between the block and the plane is 0.8. If the frictional force on the block is 10 N, the mass of the block is ($g = 10 \text{ ms}^{-2}$)

1. a. 1 kg
 2. b. 2 kg
 3. c. 3 kg
 4. d. 4 kg
-

Question 9: Two particles of masses m_1 and m_2 have equal kinetic energies. The ratio of their moments is

1. a. $m_1 : m_2$
 2. b. $m_2 : m_1$
 3. c. $\sqrt{m_1} : \sqrt{m_2}$
 4. d. $m_1^2 : m_2^2$
-

Question 10: The pressure at the bottom of a liquid tank is not proportional to the

1. a. Acceleration due to gravity
 2. b. The density of the liquid
 3. c. Height of the liquid
 4. d. Area of the liquid surface
-

Question 11: A Carnot engine takes 300 calories of heat from a source at 500 K and rejects 150 calories of heat to the sink. The temperature of the sink is

1. a. 125 K
 2. b. 250 K
 3. c. 750 K
 4. d. 1000 K
-

Question 12: Pressure of an ideal gas is increased by keeping the temperature constant. The kinetic energy of molecules

1. a. Decreases
2. b. Increases
3. c. Remains the same
4. d. Increases or decreases depending on the nature of gas

Question 13: A man weighing 60 kg is in a lift moving down with an acceleration of 1.8 ms^{-2} . The force exerted by the floor on him is

1. a. 588 N
2. b. 480 N
3. c. Zero
4. d. 696 N

Question 14: Moment of inertia of a body about two perpendicular axes X and Y in the plane of the lamina are 20 kg m^2 respectively. Its moment of inertia about an axis perpendicular to the plane of the lamina and passing through the point of intersection of X and Y axes is

1. a. 5 kg m^2
2. b. 45 kg m^2
3. c. 12.5 kg m^2
4. d. 500 kg m^2

Question 15: Two wires A and B are stretched by the same load. If the area of cross-section of wire 'A' is double that of 'B', then the stress on 'B' is

1. a. Equal to that on A
2. b. Twice that on A
3. c. Half that on A
4. d. Four times that on A

Question 16: The magnitude of point charge due to which the electric field 30 cm away has the magnitude 2 NC^{-1} will be

1. a. $2 \times 10^{-11} \text{ C}$
 2. b. $3 \times 10^{-11} \text{ C}$
 3. c. $5 \times 10^{-11} \text{ C}$
 4. d. $9 \times 10^{-11} \text{ C}$
-


Question 17: A mass of 1 kg carrying a charge of 2 C is accelerated through a potential of 1 V. The velocity acquired by it is

1. a. $\sqrt{2} \text{ ms}^{-1}$
 2. b. 2 ms^{-1}
 3. c. $1 / \sqrt{2} \text{ ms}^{-1}$
 4. d. $1 / 2 \text{ ms}^{-1}$
-

Question 18: The force of repulsion between two identical positive charges when kept, with a separation 'r' in the air is 'F'. Half the gap between the two charges is filled by a dielectric slab of dielectric constant = 4. Then, the new force of repulsion between those two charges becomes

1. a. $F / 3$
 2. b. $F / 2$
 3. c. $F / 4$
 4. d. $4F / 9$
-

Question 19: For the arrangement of capacitors as shown in the circuit, the effective capacitance between point A and B is (capacitance of each capacitor is $4\mu\text{F}$)



1. a. $4 \mu\text{F}$
 2. b. $2 \mu\text{F}$
 3. c. $1 \mu\text{F}$
 4. d. $8 \mu\text{F}$
-

Question 20: The work done to move a charge on an equipotential surface is

1. a. Infinity
2. b. Less than 1
3. c. Greater than 1
4. d. Zero

Question 21: Two capacitors of $3\ \mu\text{F}$ and $6\ \mu\text{F}$ are connected in series and a potential difference of $900\ \text{V}$ is applied across the combination. They are then disconnected and reconnected in parallel. The potential difference across the combination is

1. a. Zero
 2. b. $100\ \text{V}$
 3. c. $200\ \text{V}$
 4. d. $400\ \text{V}$
-


Question 22: Ohm's Law is applicable to

1. a. Diode
 2. b. Transistor
 3. c. Electrolyte
 4. d. Conductor
-

Question 23: If the last band on the carbon resistor is absent, then the tolerance is


1. a. $5\ \%$
 2. b. $20\ \%$
 3. c. $10\ \%$
 4. d. $15\ \%$
-

Question 24: The effective resistance between P and Q for the following network is



1. a. $(1 / 12) \Omega$
 2. b. 21Ω
 3. c. 12Ω
 4. d. $(1 / 21) \Omega$
-

Question 25: Five identical resistors each of resistance $R = 1500$ are connected to a 300 V battery as shown in the circuit. The reading of the ideal ammeter A is




1. a. $(1 / 5) \text{ A}$
 2. b. $(3 / 5) \text{ A}$
 3. c. $(2 / 5) \text{ A}$
 4. d. $(4 / 5) \text{ A}$
-

Question 26: Two cells of internal resistances r_1 and r_2 and of the same emf are connected in series, across a resistor of resistance R . If the terminal potential difference across the cells of internal resistance r_1 is zero, then the value of R is

1. a. $R = 2 (r_1 + r_2)$
 2. b. $R = r_2 - r_1$
 3. c. $R = r_1 - r_2$
 4. d. $R = 2 (r_1 - r_2)$
-

Question 27: The $I - V$ graphs for two different electrical appliances P and Q are shown in the diagram. If R_P and R_Q be the resistances of the devices, then



1. a. $R_p = R_q$
2. b. $R_p > R_q$
3. c. $R_p < R_q$
4. d. $R_p = R_q / 2$

Question 28: The correct Biot-Savart law in vector form is

1. a. $d\vec{B} = \mu_0 4\pi I (d\vec{l} \times \vec{r}) r^2$
2. b. $d\vec{B} = \mu_0 4\pi I (d\vec{l} \times \vec{r}) r^3$
3. c. $d\vec{B} = \mu_0 4\pi I (d\vec{l}) r^2$
4. d. $d\vec{B} = \mu_0 4\pi I (d\vec{l}) r^3$

Question 29: An electron is moving in a circle of radius r in a uniform magnetic field B . Suddenly, the field is reduced to $B / 2$. The radius of the circular path now becomes

1. a. $r / 2$
2. b. $2r$
3. c. $r / 4$
4. d. $4r$

Question 30: A charge q is accelerated through a potential difference V . It is then passed normally through a uniform magnetic field, where it moves in a circle of radius r . The potential difference required to move it in a circle of radius $2r$ is

1. a. $2V$
2. b. $4V$
3. c. $1V$
4. d. $3V$

Question 31: A cyclotron's oscillator frequency is 10 MHz and the operating magnetic field is 0.66 T. If the radius of its dees is 60 cm, then the kinetic energy of the proton beam produced by the accelerator is

1. a. 9 MeV
 2. b. 10 MeV
 3. c. 7 MeV
 4. d. 11 MeV
-

Question 32: Needles N_1 , N_2 and N_3 are made of a ferromagnetic, a paramagnetic and a diamagnetic substance respectively. A magnet, when brought close to them, will

1. a. Attract all three of them
 2. b. Attract N_1 strongly, N_2 weakly and repel N_3 weakly
 3. c. Attract N_1 strongly but repel N_2 and N_3 weakly
 4. d. Attract N_1 and N_2 strongly but repel N_3
-





Question 33: The strength of the Earth's magnetic field is

1. a. Constant everywhere
 2. b. Zero everywhere
 3. c. Having very high value
 4. d. Varying from place to place on the Earth's surface
-

Question 34: A jet plane with a wing-span of 25 m is travelling horizontally towards the east with a speed of 3600 km/hour. If the Earth's magnetic field at the location is 4×10^{-4} T and the angle of dip is 30° , then, the potential difference between the ends of the wing is

1. a. 4 V
 2. b. 5 V
 3. c. 2 V
 4. d. 2.5 V
-

Question 35: Which of the following represents the variation of inductive reactance (X_L) with the frequency of voltage source (ν)?



Question 36: The magnetic flux linked with a coil varies as $\Phi = 3t^2 + 4t + 9$. The magnitude of the emf induced at $t = 2$ seconds is

1. a. 8 V
2. b. 16 V
3. c. 32 V
4. d. 64 V

Question 37: A 100 W bulb is connected to an AC source of 220 V, 50 Hz. Then the current flowing through the bulb is

1. a. $[5 / 11]$ A
2. b. $[1 / 2]$ A
3. c. 2 A
4. d. $[3 / 4]$ A

Question 38: In the series LCR circuit, the power dissipation is through

1. a. R
2. b. L
3. c. C
4. d. Both L and C

Question 39: In Karnataka, the normal domestic power supply AC is 220 V, 50 Hz. Here 220 V and 50 Hz refer to

1. a. Peak value of voltage and frequency
2. b. Rms value of voltage and frequency
3. c. Mean value of voltage and frequency
4. d. Peak value of voltage and angular frequency

Question 40: A step-up transformer operates on a 230 V line and I loads current of 2 A. The ratio of primary and secondary windings is 1:25. Then the current in the primary is

1. a. 25 A
2. b. 50 A
3. c. 15 A
4. d. 12.5 A

Question 41: The number of photons falling per second on a completely darkened plate to produce a force of 6.62×10^{-5} N is 'n'. If the wavelength of the light falling is 5×10^{-7} m, then $n = \text{_____} \times 10^{22}$. ($h = 6.62 \times 10^{-34}$ J-s)

1. a. 1
2. b. 5
3. c. 0.2
4. d. 3.3

Question 42: An object is placed at the principal focus of a convex mirror. The image will be at

1. a. Centre of curvature
2. b. Principal focus
3. c. Infinity
4. d. No image will be formed

Question 43: An object is placed at a distance of 20 cm from the pole of a concave mirror of focal length 10 cm. The distance of the image formed is

1. a. + 20 cm
 2. b. + 10 cm
 3. c. -20 cm
 4. d. - 10 cm
-

Question 44: A candle placed 25 cm from a lens forms an image on the screen placed 75 cm on the other side of the lens. The focal length and type of the lens should be

1. a. + 18.75 cm and convex lens
 2. b. - 18.75 cm and concave lens
 3. c. + 20.25 cm and convex lens
 4. d. -20.25 cm and concave lens
-

Question 45: A plane wavefront of wavelength λ is incident on a single slit of width a . The angular width of the principal maximum is

1. a. λ / a
 2. b. $2\lambda / a$
 3. c. a / λ
 4. d. $a / 2\lambda$
-

Question 46: In a Fraunhofer diffraction at a single slit, if yellow light illuminating the slit is replaced by blue light, then diffraction bands

1. a. Remain unchanged
 2. b. Become wider
 3. c. Disappear
 4. d. Become narrower
-

Question 47: In Young's double-slit experiment, two wavelengths $\lambda_1 = 780 \text{ nm}$ and $\lambda_2 = 520 \text{ nm}$ are used to obtain interference fringes. If the n^{th} bright band due to λ_1 coincides with $(n + 1)^{\text{th}}$ bright band due to λ_2 , then the value of n is

1. a. 4
 2. b. 3
 3. c. 2
 4. d. 6
-

Question 48: In Young's double-slit experiment, slits are separated by 2 mm and the screen is placed at a distance of 1.2 m from the slits. Light consisting of two wavelengths 6500 \AA and 5200 \AA is used to obtain interference fringes. Then the separation between the fourth bright fringes of two different patterns produced by the two wavelengths is

1. a. 0.312 mm
 2. b. 0.123 mm
 3. c. 0.213 mm
 4. d. 0.412 mm
-

Question 49: The maximum kinetic energy of emitted photoelectrons depends on

1. a. Intensity of incident radiation
 2. b. Frequency of incident radiation
 3. c. Speed of incident radiation
 4. d. Number of photons in the incident radiation
-

Question 50: A proton and an α particle are accelerated through the same potential difference V . The ratio of their de Broglie wavelengths is

1. a. $\sqrt{2}$
2. b. $2\sqrt{2}$
3. c. $\sqrt{3}$
4. d. $2\sqrt{3}$

Question 51: The total energy of an electron revolving in the second orbit of the hydrogen atom is

1. a. -13.6 eV
2. b. -1.51 eV
3. c. -3.4 eV
4. d. Zero

Question 52: The period of revolution of an electron in the ground state of the hydrogen atom is T . The period of revolution of the electron in the first excited state is

1. a. $2T$
2. b. $4T$
3. c. $6T$
4. d. $8T$

Question 53: The energy equivalent to a substance of mass 1 g is

1. a. 18×10^{13} J
2. b. 9×10^{13} J
3. c. 18×10^6 J
4. d. 9×10^6 J

Question 54: The half-life of tritium is 12.5 years. What mass of tritium of initial mass 64 mg will remain undecayed after 50 years?

1. a. 32 mg
2. b. 8 mg
3. c. 16 mg
4. d. 4 mg

Question 55: In a CE amplifier, the input ac signal to be amplified is applied across

1. a. Forward biased emitter-base junction
2. b. Reverse biased collector-base junction
3. c. Reverse biased emitter-base junction
4. d. Forward biased collector-base junction

Question 56: If $A = 1$ and $B = 0$, then in terms of Boolean algebra, $A^{-} + B =$

1. a. B
 2. b. B^{-}
 3. c. A
 4. d. A^{-}
-

Question 57: The density of an electron-hole pair in pure germanium is $3 \times 10^{16} \text{ m}^{-3}$ at room temperature. On doping with aluminium, the hole density increases to $4.5 \times 10^{22} \text{ m}^{-3}$. Now the electron density (in m^{-3}) in doped germanium will be

1. a. 1×10^{10}
 2. b. 2×10^{10}
 3. c. 0.5×10^{10}
 4. d. 4×10^{10}
-

Question 58: The dc common-emitter current gain of a n-p-n transistor is 50. The potential difference applied across the collector and emitter of a transistor used in CE configuration is, $V_{CE} = 2 \text{ V}$. If the collector resistance, $R_C = 4 \Omega \text{ k}$, the base current (I_B) and the collector current (I_C) are

1. a. $I_B = 10 \mu\text{A}$, $I_C = 0.5 \text{ mA}$
 2. b. $I_B = 0.5 \mu\text{A}$, $I_C = 10 \text{ mA}$
 3. c. $I_B = 5 \mu\text{A}$, $I_C = 1 \text{ mA}$
 4. d. $I_B = 1 \mu\text{A}$, $I_C = 0.5 \text{ mA}$
-

Question 59: The radius of the Earth is 6400 km. If the height of an antenna is 500 m, then its range is

1. a. 800 km
2. b. 100 km
3. c. 80 km
4. d. 10 km

Question 60: A space station is at a height equal to the radius of the Earth. If ' V_E ' is the escape velocity on the surface of the Earth, the same on the space station is _____ times V_E .

1. a. $1/2$
2. b. $1/4$
3. c. $1/\sqrt{2}$
4. d. $1/\sqrt{3}$