



**KLS**

**Vishwanathrao Deshpande Institute of Technology  
Haliyal - 581329**

---

**FIRST YEAR - CHEMISTRY CYCLE**

**AIML**

**LAB Manual**

**2025 Scheme**

NAME : \_\_\_\_\_  
Roll.No./ USN : \_\_\_\_\_  
DIVISION : \_\_\_\_\_

<b>Semester : II</b>			
<b>Sl.No.</b>	<b>Lab Details</b>	<b>Lab Code</b>	<b>Pg. No.</b>
1	ENGINEERING CHEMISTRY	1BCHES102/202	04-65
2	PYTHON PROGRAMMING	1BPLC105B/205B	66-85



As prescribed by,  
**VISVESVARAYA TECHNOLOGICAL UNIVERSITY,  
BELAGAVI - 590014**

## **PROGRAM OUTCOMES (POs)**

Program Outcomes as defined by NBA (PO) Engineering Graduates will be able to:

1. **PO1: Engineering Knowledge:** Apply knowledge of mathematics, natural science, computing, engineering fundamentals and an engineering specialization as specified in WK1 to WK4 respectively to develop to the solution of complex engineering problems.
2. **PO2: Problem Analysis:** Identify, formulate, review research literature and analyze complex engineering problems reaching substantiated conclusions with consideration for sustainable development. (WK1 to WK4)
3. **PO3: Design/Development of Solutions:** Design creative solutions for complex engineering problems and design/develop systems/components/processes to meet identified needs with consideration for the public health and safety, whole-life cost, net zero carbon, culture, society and environment as required. (WK5)
4. **PO4: Conduct Investigations of Complex Problems:** Conduct investigations of complex engineering problems using research-based knowledge including design of experiments, modeling, analysis & interpretation of data to provide valid conclusions. (WK8).
5. **PO5: Engineering Tool Usage:** Create, select and apply appropriate techniques, resources and modern engineering & IT tools, including prediction and modeling recognizing their limitations to solve complex engineering problems. (WK2 and WK6)
6. **PO6: The Engineer and The World:** Analyze and evaluate societal and environmental aspects while solving complex engineering problems for its impact on sustainability with reference to economy, health, safety, legal framework, culture and environment. (WK1, WK5, and WK7).
7. **PO7: Ethics:** Apply ethical principles and commit to professional ethics, human values, diversity and inclusion; adhere to national & international laws. (WK9)
8. **PO8: Individual and Collaborative Team work:** Function effectively as an individual, and as a member or leader in diverse/multi-disciplinary teams.
9. **PO9: Communication:** Communicate effectively and inclusively within the engineering community and society at large, such as being able to comprehend and write effective reports and design documentation, make effective presentations considering cultural, language, and learning differences
10. **PO10: Project Management and Finance:** Apply knowledge and understanding of engineering management principles and economic decision-making and apply these to one's own work, as a member and leader in a team, and to manage projects and in multidisciplinary environments.
11. **PO11: Life-Long Learning:** Recognize the need for, and have the preparation and ability for i) independent and life-long learning ii) adaptability to new and emerging technologies and iii) critical thinking in the broadest context of technological change. (WK8)

### **COLLEGE VISION AND MISSION**

<b>Vision ( College )</b>
To nurture talent & enrich society through excellence in technical education, research & innovation.
<b>Mission ( College )</b>
<ol style="list-style-type: none"> <li>1. To augment innovative Pedagogy, &amp; kindle quest for interdisciplinary learning &amp; to enhance conceptual understanding.</li> <li>2. To build competence, professional ethics &amp; develop entrepreneurial thinking.</li> <li>3. To strengthen Industry Institute Partnership &amp; explore global collaborations.</li> <li>4. To inculcate culture of socially responsible citizenship.</li> <li>5. To focus on Holistic &amp; Sustainable development</li> </ol>

<b>Vision ( chemistry dept )</b>
To grow profusely as departments of par excellence in the area of scientific education developing human resources of towering talents with sensible temperament.
<b>Mission ( chemistry dept )</b>
To train students to emerge as exceptionally skilled technocrats enthralled with scientific approach, with a sense of social responsibility.
<b>Vision ( CSE AI&amp;ML Dept )</b>
To lead the way in the creation and application of cutting-edge Computer science and engineering (AIML) technologies, advancing the frontiers of knowledge, and empowering future generations to drive innovation and transformation on a global scale.
<b>Mission ( CSE AI&amp;ML Dept )</b>
<p>To train students with a strong conceptual understanding using innovative pedagogies, empowering them to excel in the dynamic fields of Artificial Intelligence and Machine Learning.</p> <p>To imbibe professional, research, and entrepreneurial skills with a commitment to the nation's development at large.</p> <p>To strengthen the industry-institute Interaction.</p> <p>To promote life-long learning with a sense of societal &amp; ethical responsibilities.</p>

# LAB MANUAL-1

## ENGINEERING CHEMISTRY

(1BCHE102/202)

### CO's and PO's Mapping Chart

Sl. No.	Description	PO 1	PO2	PO3	PO4	PO5	PO6	PO7	PO8	PO9	PO 10	PO 11
1	Understand and analyze the properties, classification and applications of semiconductor materials, energy storage and conversion devices.	3	2	1	1		1					1
2	Demonstrate knowledge of nanomaterials and quantum dots including their synthesis, properties, and device applications.	3	2	1			1					1
3	Explain the role of functional polymers and composites in flexible electronic applications.	3	2	1					1			1
4	Apply experimental skills and electrochemical concepts to sensor systems and evaluate corrosion control and e-waste management techniques	3	2	1	1					1		1

**Evaluation:**

<b>Integrated Lab(IPCC)</b>		
<b>CIE</b>		
<b>Particulars</b>	<b>Marks</b>	<b>Total</b>
Conduction	04	
Performance	03	
Journal	02	
Viva Voce	01	
	<b>10</b>	
<b>Total Marks</b>	12x10	<b>120</b>
	Reduced	<b>15</b>
<b>Lab IA</b>		
<b>Particulars</b>	<b>Marks</b>	<b>Total</b>
IA	<b>50</b>	10 (Reduced)
Procedure write Up- 08 Conduction- 26 Observation and Calculation- 09 Viva- 07		
<b>Total (CIE +Lab IA)</b>		<b>25</b>

### Mapping of Experiments with CO and PO

S.No.	Experiment Details	CO	PO
1	Estimation of total hardness of water by EDTA method	1,2,3,4	1,2,3,4,6,8,9,11
2	Determination of chemical oxygen demand (COD) of industrial effluent sample.	1,2,3,4	1,2,3,4,6,8,9,11
3	Estimation of iron in TMT bar by diphenyl amine indicator method.	1,2,3,4	1,2,3,4,6,8,9,11
4	Determination of total alkalinity of given water sample.	1,2,3,4	1,2,3,4,6,8,9,11
5	Determination of acid value of solder flux using KOH	1,2,3,4	1,2,3,4,6,8,9,11
6	Estimation of acid mixture by conductometric sensor (Conductometry)	1,2,3,4	1,2,3,4,6,8,9,11
7	Estimation of iron in rust sample by Potentiometric sensor (Potentiometry)	1,2,3,4	1,2,3,4,6,8,9,11
8	Determination of pKa of vinegar using pH sensor (Glass electrode)	1,2,3,4	1,2,3,4,6,8,9,11
9	Estimation of copper present in e-waste by optical sensor (Colorimetry).	1,2,3,4	1,2,3,4,6,8,9,11
10	Determination of Viscosity coefficient of Coolant Samples	1,2,3,4	1,2,3,4,6,8,9,11
11	Interpretation of pka values of a cooling fluid using origin software.	1,2,3,4	1,2,3,4,6,8,9,11
12	Chemical structure drawing using software: Chem Draw/ Chem Sketch.	1,2,3,4	1,2,3,4,6,8,9,11

**EXPERIMENT WISE LESSON PLAN**

<b>Experiment No.1</b>	
<b>Name</b>	Estimation of total hardness of water by EDTA method
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To enable students to understand the concept of water hardness and its significance.</li> <li>To develop skills in accurate titration and endpoint detection.</li> </ul>
<b>Experiment No.2</b>	
<b>Name</b>	Determination of chemical oxygen demand (COD) of industrial effluent sample
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To help students understand the importance of COD in environmental monitoring.</li> <li>To familiarize students with oxidation–reduction titration methods.</li> <li>To develop awareness about industrial wastewater pollution.</li> </ul>
<b>Experiment No.3</b>	
<b>Name</b>	Estimation of iron in TMT bar by diphenyl amine indicator method.
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To enable students to determine metal content in engineering materials.</li> <li>To enhance analytical skills in quantitative chemical analysis.</li> </ul>
<b>Experiment No.4</b>	
<b>Name</b>	Determination of total alkalinity of given water sample.
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To help students understand alkalinity and its role in water quality.</li> <li>To train students in acid–base titration techniques.</li> </ul>
<b>Experiment No. 5</b>	
<b>Name</b>	Determination of acid value of solder flux using KOH

<b>Objectives</b>	<ul style="list-style-type: none"> <li>To familiarize students with the analysis of industrial chemical products.</li> <li>To improve students' laboratory handling and safety skills.</li> </ul>
<b>Experiment No. 6</b>	
<b>Name</b>	Estimation of acid mixture by conductometric sensor (Conductometry)
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To help students understand the relationship between conductivity and concentration.</li> <li>To develop skills in using modern analytical instruments.</li> </ul>
<b>Experiment No. 7</b>	
<b>Name</b>	Estimation of iron in rust sample by Potentiometric sensor (Potentiometry)
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To help students understand corrosion analysis.</li> <li>To develop competence in handling electrochemical sensors.</li> </ul>
<b>Experiment No. 8</b>	
<b>Name</b>	Determination of pKa of vinegar using pH sensor (Glass electrode)
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To help students understand acid dissociation and pKa concepts.</li> <li>To train students in the use of pH meters.</li> <li>To develop skills in interpreting titration curves.</li> </ul>
<b>Experiment No.9</b>	
<b>Name</b>	Estimation of copper present in e-waste by optical sensor (Colorimetry).
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To introduce students to colorimetric analysis techniques.</li> <li>To raise awareness about e-waste recycling and resource recovery.</li> <li>To develop skills in optical measurement and data analysis</li> </ul>
<b>Experiment No.10</b>	
<b>Name</b>	Determination of Viscosity coefficient of Coolant Samples
<b>Objectives</b>	<ul style="list-style-type: none"> <li>To help students understand viscosity and fluid properties.</li> <li>To provide practical exposure to rheological measurements.</li> <li>To relate material properties to engineering applications.</li> </ul>

---

<b>Experiment No.11</b>	
<b>Name</b>	Interpretation of pka values of a cooling fluid using origin software.
<b>Objectives</b>	<ul style="list-style-type: none"><li>• To expose students to modern, low-cost analytical techniques.</li><li>• To demonstrate applications of chemistry in pH analysis.</li><li>• To develop skills in digital data acquisition and analysis.</li></ul>
<b>Experiment No.12</b>	
<b>Name</b>	Chemical structure drawing using software: Chem Draw/ Chem Sketch.
<b>Objectives</b>	<ul style="list-style-type: none"><li>• To introduce students to draw chemical structures .</li><li>• To easily interpret the chemical compounds.</li></ul>

**LIST OF EXPERIMENTS**

S. No.	Experiment	Page. No
1	Estimation of total hardness of water by EDTA method	12-17
2	Determination of chemical oxygen demand (COD) of industrial effluent sample.	18-24
3	Estimation of iron in TMT bar by diphenyl amine indicator method.	25-28
4	Determination of total alkalinity of given water sample.	29-32
5	Determination of acid value of solder flux using KOH	33-35
6	Estimation of acid mixture by conductometric sensor (Conductometry)	36-42
7	Estimation of iron in rust sample by Potentiometric sensor (Potentiometry).	43-48
8	Determination of pKa of vinegar using pH sensor (Glass electrode)	49-51
9	Estimation of copper present in e-waste by optical sensor (Colorimetry).	52-56
10	Determination of Viscosity coefficient of Coolant Samples	57-60
11	Interpretation of pka values of a cooling fluid using origin software	61-62
12	Chemical structure drawing using software: Chem Draw/ Chem Sketch.	63-65

### **LAB SAFETY & USAGE INSTRUCTIONS**

1. Study theory behind the experiment before attending the Laboratory.
2. Keep the work bench and sink (wash basin) neat and clean. Do not allow used filter papers, broken pieces of glass, used match sticks, etc., to lie on the work bench –throw them into the available dust bin nearby.
3. Keep the apparatus clean and arrange them properly.
4. Handle the chemicals and reagent bottles carefully.
5. Take the prescribed quantities of chemicals and reagents only.
6. Do not pour any excess reagent, taken by chance, back into the reagent bottle, as it is likely to contaminate the entire solution in the reagent bottle.
7. Close the reagent bottles with their lids and keep them in their proper places, after use.
8. Water is a precious commodity; do not waste it; close the water tap immediately after use.
9. It is said, 'Prevention is better than cure' - take care to prevent fire accidents in the Lab.
10. Make it a habit to record all observations in your Observation Note Book, as and when you carry out an experiment; writing observations on loose bits of paper is a bad habit.
11. Do not forget to bring your Laboratory Record while attending the lab.
12. Always wear shoes and laboratory apron while you are in the lab.
13. Wash chemical spills on your body, if any, immediately with plenty of tap water.
14. Before leaving the Laboratory, wash the apparatus clean, keep them in proper place and make your work bench tidy.

## Experiment 1

### Estimation of Total Hardness of Water by EDTA Method

**Aim:** To determine the total hardness of water

**Chemicals:** Na<sub>2</sub>EDTA, Ammonia, NH<sub>4</sub>OH-NH<sub>4</sub>Cl buffer & Eriochrome black-T indicator

**Theory:** Hardness of water is the soap-consuming factor of water. Hard water does not lather well with soap. Soft water lathers well with soap. There are two types in hardness. Temporary hardness (can be removed by simply boiling the water) and permanent hardness (cannot be removed by boiling instead, can be removed by chemical methods). Hard water poses problems of excess soap consumption in domestic regions and scale & sludge formations in boilers and in industries. Soft water is most suited for many applications. Bicarbonates of calcium and magnesium cause temporary hardness. Chlorides, sulfates & nitrates of calcium and magnesium cause permanent hardness. It may be noted that hardness is due to Ca<sup>2+</sup> & Mg<sup>2+</sup>

ions and not due to the anions such as, Cl<sup>-</sup>, NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, HCO<sup>-</sup>, etc. Total hardness is the sum of temporary hardness and permanent hardness. Though, hardness is due to different metal salts, it is expressed in equivalent amounts of CaCO<sub>3</sub> (CaCO<sub>3</sub> is chosen because, it is most insoluble in water and its molecular weight is a rounded number, 100). Hardness is expressed in terms of ppm (parts of CaCO<sub>3</sub> equivalents per million parts of water) or mg/L (milligrams of CaCO<sub>3</sub> equivalents per liter of water).

Hardness is determined by EDTA complexometric method (EDTA = Ethylene Diamine Tetra Acetic acid). The method involves the titration of known volume of hard water against standard EDTA solution using eriochrome black-T (EBT) as metal ion indicator at pH = 10 (reaction is complete at this pH and colour change happens to be sharp at the end point). The indicator forms relatively less stable wine red coloured Ca-EBT & Mg-EBT complexes. When EDTA solution is added, initially it will react with freely available Mg<sup>2+</sup> & Ca<sup>2+</sup> ions and towards the end point, it snatches Ca<sup>2+</sup> and Mg<sup>2+</sup> ions from the less stable Ca-EBT & Mg-EBT complexes to form more stable, colourless Ca-EDTA & Mg-EDTA complexes. Free eriochrome black-T is liberated which

imparts blue colour in the medium of pH =10. This marks the end of titration.

On adding indicator:  $\text{Ca}^{2+} + \text{EBT} \longrightarrow \text{Ca-EBT complex}$

$\text{Mg}^{2+} + \text{EBT} \longrightarrow \text{Mg-EBT complex}$   
(wine red)

At end point:  $\text{Ca-EBT complex} + \text{EDTA} \longrightarrow \text{Ca-EDTA complex} + \text{EBT}$   
(wine red) (colourless) (blue)

$\text{Mg-EBT complex} + \text{EDTA} \longrightarrow \text{Mg-EDTA complex} + \text{EBT}$   
(wine red) (colourless) (blue)

Knowing the molarity and volume of EDTA consumed, hardness of water is calculated.

### Procedure:

#### A. Preparation of standard $\text{Na}_2\text{EDTA}$ solution:

Take the beaker containing about 1.0 g  $\text{Na}_2\text{EDTA}$  crystals and note down the beaker number. Add 5 mL of 1:1 ammonia solution, a little of distilled water and dissolve the crystals. Transfer the solution into a 250 mL capacity standard flask through a funnel. Rinse the beaker with little of distilled water at a time and transfer the rinsing into the standard flask. Dilute the solution up to the mark with distilled water. Shake well to make the solution uniform.

#### B. Determination of total hardness of water:

Take a clean burette and rinse it with distilled water and then with standard EDTA solution prepared. Now, fill the burette with standard EDTA solution. Take a clean conical flask and rinse it with distilled water. Take a clean pipette and rinse it with the given sample of hard water. Pipette out 25 mL of hard water into the conical flask. Add 2 mL of buffer solution ( $\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$ ) and a pinch or 2-3 drops of eriochrome black-T indicator (it is also known as, solochrome black). Titrate this solution against standard  $\text{Na}_2\text{EDTA}$  solution adding 1.0 mL at a time until the change in colour from wine red to clear blue (pilot reading). Repeat the titration adding 0.1 mL towards the end point for two correct readings. Let the average of burette readings be P mL. Calculate the total hardness of the given sample of water.

#### Result:

The total hardness of the given sample of water= ..... ppm (or mg/L)

**Observations and calculations:****A. Preparation of standard Na<sub>2</sub>EDTA solution:**

Beaker number = .....

Weight of Na<sub>2</sub>EDTA taken (*given*) = W = .....gVolume of standard solution prepared from W g of Na<sub>2</sub>EDTA = 250 mlMolarity of Na<sub>2</sub>EDTA solution, Y =  $\frac{4 \times W}{372.25} = \frac{4 \times \dots\dots\dots}{372.25} = \dots\dots\dots$ M**B. Determination of total hardness of water:**Solution taken in burette : Y = ..... molar Na<sub>2</sub>EDTA solutionSolution taken in conical flask : 25 mL of hard water sample  
+ 2 mL NH<sub>4</sub>OH-NH<sub>4</sub>Cl buffer solution (pH = 10)

Indicator : Eriochrome black – T (a pinch of indicator powder or 2 drops of solution)

Color change at end point : Wine red to clear blue

**Burette readings:**

	Trial-1 (Pilot)	Trial-2	Trial-3
Final Reading			
Initial Reading			
Difference (ml)			

Average of agreeing volumes (P) = .....ml

It is known that

1000 mL of 1 molar Na<sub>2</sub>EDTA solution = 100 g of CaCO<sub>3</sub>(Molecular weight of CaCO<sub>3</sub> = 100)1 mL of 1 molar Na<sub>2</sub>EDTA solution = 0.1 g of CaCO<sub>3</sub>

$$\begin{aligned}
 P \text{ mL of } Y \text{ molar Na}_2\text{EDTA solution} &= 0.1 \times P \times Y \text{ g of CaCO}_3 \\
 &= 0.1 \times \dots \times \dots \text{ g of} \\
 &\quad \text{CaCO}_3
 \end{aligned}$$

$$Z = \dots \text{ g of CaCO}_3$$

$$\text{Now, 25 mL (= 25 g) of hard water} = Z = \dots \text{ g of CaCO}_3$$

$$\begin{aligned}
 \therefore 10^6 \text{ mL } (\cong 10^6 \text{ g}) \text{ of hard water} &= \frac{ZX10^6}{25} = \frac{\dots \times 10^6}{25} \\
 &= \dots \text{ g of CaCO}_3
 \end{aligned}$$

Hence, total hardness of given sample of water

$$= \dots \text{ grams per million grams}$$

$$= \dots \text{ ppm or mg/ml}$$

### VIVA QUESTIONS & ANSWERS

**1. What is meant by hard water?**

Ans. Water not giving lather with soap is known as hard water.

**2. What is the unit of hardness of water?**

Ans. It is ppm of calcium carbonate.


**3. What are the types of hard water?**

Ans. Temporary due to the presence of bicarbonates of calcium and magnesium  
Permanent hard ness is due to the presence chlorides and sulfates of calcium and magnesium.

**4. What is the type of titration is carried out for the estimation of total hardness?**

Ans: Complexometric titration with EDTA salt

**5. What is the structural formula of EDTA?**



**6. Why disodium salt of EDTA is preferred to EDTA?**

Ans. Disodium salt dissolves readily in water, whereas EDTA is sparingly soluble

**7. How do you calculate Molarity of EDTA?**

Ans. Molarity of EDTA = Number of grams of EDTA in 1000 ml / Molecular wt. (372)

**8. What is a buffer solution?**

Ans. Addition small quantity of acid or alkali does not alter the pH of solution.

**9. Why ammonia solution is added while preparing standard solution of EDTA?**

Ans. This is to increase the rate of dissolution of EDTA in water.

**10. Why is ammonia-ammonium chloride buffer added?**

Ans. This buffer is added to maintain pH of 10-11

**11. Why is the indicator Eriochrome Black T shows wine red at the beginning and blue color at the end?**

Ans. The indicator forms weak complex with calcium and magnesium present in hard water and gives wine red color at pH 10. During titration the EDTA forms stronger complex with calcium and magnesium ions and at the end point EDTA liberates the indicator, which is blue in color at pH 10.

**12. Why is titration-involving EDTA carried out slowly towards end point?**

Ans. Since the formation of metal complex is very slow, the titration has to be done slowly.

**13. What is importance of hardness data?**

Ans. It is important in determining the suitability of water for domestic and industrial use. Determination of hardness of water serves as a basis for routine control of softening process.

**14. Define ppm?**

Ans: No. of parts of  $\text{CaCO}_3$  equivalent to hardness causing ions present in million parts of water.

**15. Why is hardness expressed only in terms of  $\text{CaCO}_3$ ?**

Ans : Because it is highly insoluble salt and its molecular weight is 100 which make the calculation easier.

**16. How many bonding sites are present in EDTA?**

Ans : Six bonding sites (called hexadentate ligand). Namely, two nitrogen atoms and four oxygen atoms.

**17. What is meant by ligand?**

Ans : A species which can donate a lone pair of electrons to the metal atom is called ligand.

**18. What is buffer solution?**

Ans : Solution which resist the change in the pH of the solution upon the

addition of small amount of acid or base.

**19. How pH change takes place during the course of the reaction?**

Ans :Metal ion combines with EDTA to form complex, leading to the liberation of  $H^+$  ions which results in the change in pH of the solution.

**20. Give the other name for Eriochrome black-T indicator?**

Ans :Solochrome black-T.

**21. What is the end point in the determination of total hardness of water?**

Ans :Change of color from wine red to pure blue.

**22. How does water acquire temporary hardness?**

Ans : Due to the presence of bicarbonates of calcium and magnesium ions.

**23. How do you remove temporary hardness?**

Ans : By boiling the water sample.

**24. What happens when the hard water sample is boiled?**

Ans :The bicarbonates salts of calcium and magnesium get converted into insoluble carbonates. This can be removed by decantation or filtration.

**25. How are the waters classified based on the degree of hardness?**

Ans : Soft water	0 – 75 mg/litre
Moderately hard water	75 – 150 mg/litre
Hard water	150 – 300 mg/litre
Very hard water	> 300 mg/litre

**26. What do you mean by total hardness of water?**

Ans : The sum of both temporary and permanent hardness is called total hardness of water.



**Procedure:****A Preparation of standard FAS solution:**

Take the beaker containing about 5-7 g of FAS and note down the beaker number. Add 1 test tube full of dilute  $\text{H}_2\text{SO}_4$  followed by about 50 mL distilled water, dissolve and transfer the solution into a 250 mL standard flask. Wash the beaker and transfer the rinsing also into the standard flask. Dilute up to the mark with distilled water. Shake the solution well to make it uniform.

**B Determination of COD:**

Take a clean burette and rinse it with standard FAS solution. Now, fill the burette with standard FAS solution. Take a clean pipette and rinse it with the waste water sample. Pipette out 25 mL of waste water into a clean conical flask\*. Add one test tube of 1:1 dilute  $\text{H}_2\text{SO}_4$ . Pipette out 25 mL of the  $\text{K}_2\text{Cr}_2\text{O}_7$  solution in the same conical flask and add 3-4 drops of ferroin indicator. Back titrate the unreacted  $\text{K}_2\text{Cr}_2\text{O}_7$  in the conical flask against standard FAS solution from the burette in increments of 1.0 mL until change in colour from bluish green to reddish brown. Repeat the titration adding 0.1 mL towards the end point for two correct readings. Let the average of burette readings be P mL.

**C Blank titration:**

Carry out a titration similar to the main titration, taking only  $\text{K}_2\text{Cr}_2\text{O}_7$  &  $\text{H}_2\text{SO}_4$  without the sample waste water (pilot reading). Repeat the titration for two correct readings. Let the average of correct readings be Q mL. Calculate the chemical oxygen demand.

\*The sample waste water given here contains only easily oxidisable pollutants. Therefore,

- (i) mercuric sulphate &
- (ii)  $\text{Ag}^+$  ion catalyst are not added and
- (iii) refluxing the reaction mixture is avoided

**Result:**

Chemical oxygen demand (COD) of given waste water sample = .....ppm.

**Observations and calculations:****A. Preparation of standard FAS solution:**

Beaker number = .....

Weight of FAS taken for analysis (ask the staff in-charge) = W = .....g

Volume of standard solution prepared from W g of FAS = 250 mL

Normality of FAS solution,  $Y = \frac{4 \times W}{392.13} = \frac{4 \times \dots}{392.13} = \dots$ 

(Equivalent weight of FAS = Molecular weight of FAS = 392.13)

**B. Determination of COD of waste water:**

Solution taken in the burette : Y = ..... Normal FAS solution

Solution taken in the conical flask : 25 mL of waste water sample

+ 25 mL of  $K_2Cr_2O_7$  solution+ 1 test tube 1:1  $H_2SO_4$ 

Indicator : Ferroin (3-4 drops)

Color change at end : Bluish green to reddish brown

**Burette readings:**

	Trial-1 (Pilot)	Trial-2	Trial-3
Final Reading			
Initial Reading			
Difference (ml)			

Average of agreeing volumes (P) = .....ml

**C. Blank titre value (given)**Burette reading when 25 mL of acidified  $K_2Cr_2O_7$  is titrated against

Y = ..... Normal FAS solution

Q = .....ml

Now,

Consumption of  $K_2Cr_2O_7$  by

25 mL waste water in terms of FAS =  $(Q - P)$  ml of Y normal FAS

= ..... - ..... mL of ..... normal FAS.

It is known that,

1000 mL of 1 normal FAS = 8 g of oxygen

**(Equivalent weight of Oxygen = Atomic weight of Oxygen = 8)**

1 mL of 1 normal FAS = 0.008 g of oxygen

$(Q - P)$  mL of Y normal FAS =  $0.008 \times (Q - P) \times Y$  g of oxygen

=  $0.008 \times \dots \times \dots$  g of oxygen

= Z = ..... g of oxygen

Now, 25 mL of waste water requires, Z = ..... g of oxygen

10 mL of waste water requires

$$\frac{Z \times 10^6}{25}$$

-

$$\frac{\dots \times 10^6}{25}$$

= .....g of Oxygen

The chemical oxygen demand (COD)

of the given sample of waste water = .....ppm

### **VIVA QUESTIONS & ANSWERS**

#### **1. What is COD?**

Ans. It is the amount of oxygen required for chemical oxidation of organic and inorganic matter present in wastewater by a strong oxidizing agent. It is expressed mg per 1000 ml of wastewater.

**2. What is BOD?**

Ans. It is the amount of oxygen required by the microorganisms for biological oxidation of organic matter in wastewater in five days at 20°C at aerobic condition.

**3. What are oxidisable constituents of wastewater?**

Ans. The oxidisable constituents of wastewater include straight chain aliphatic compounds, aromatic compounds, alcohols, and other oxidisable materials.

**4. What general groups of organic compounds are not oxidized in the COD test?**

Ans. Aromatic hydrocarbons and pyridine are not oxidized in COD

**5. What is the role of silver sulfate?**

Ans. Silver sulfate acts as a catalyst in oxidation of straight chain aliphatic hydrocarbons and acetic acid.

**6. What is the role of mercuric sulfate?**

Ans. Chloride ions are present in wastewater and the same gives erroneous result. Mercuric ions bind the halide ions present in wastewater to form poorly ionized mercuric halide and prevent the reaction between silver ions and halide.

**7. What are the products formed in COD analysis?**

Ans. During COD analysis, organic matter is completely oxidized to carbon dioxide and water.

**8. Why is sulfuric acid added during the preparation of standard FAS solution?**

Ans. The sulfuric acid is added to prevent the hydrolysis of ferrous sulfate.

**9. What is the indicator used?**

Ans. Ferroin. It is ferrous 1, 10-phenanthroline sulfate.

**10. What is the unit of COD?**

Ans. COD is expressed as mg of oxygen per 1000 ml of wastewater.

**11. Mention a few applications of COD.**

Ans. It is extensively used in the analysis of industrial wastes

It gives an idea of substances resistant for oxidation by microorganisms

**12. What is the role of 1:1 sulfuric acid added in the conical flask?**

Potassium dichromate acts as oxidizing agent in acid medium

**13. How is 150 ml of 6N sulfuric acid prepared with 36N sulfuric acid?**

$$N_1 \times V_1 = N_2 \times V_2$$

$$6 \times 150 = 36 \times$$

$$V_2$$

$$V_2 = 6 \times 150 / 36 = 25 \text{ ml}$$

25 ml of 36N sulfuric acid is added to 125 ml of water to get 6N of 150 ml of sulfuric acid.

**14. Which is the oxidizing agent used in COD experiment?**

Acidified potassium dichromate solution.

**15. What is the full form of FAS? Give its composition?**

Ferrous ammonium sulphate.  $\text{FeSO}_4 (\text{NH}_4)_2 \text{SO}_4 \cdot 6\text{H}_2\text{O}$

**16. Why is sulphuric acid added during the preparation of std. FAS solution?**

Sulphuric acid is added to prevent the hydrolysis of ferrous sulphate to ferrous hydroxide.

**17. What is the end point in COD estimation?**

Bluish green to reddish brown.

**18. Why is COD value is higher than BOD for the same water sample?**

This is because COD involves complete oxidation of both organic and inorganic compounds, whereas BOD involves the oxidation of only the organic compounds.

**19. What is the advantage of COD over BOD?**

COD can be obtained in a short interval of time and it helps in estimation of both organic and inorganic oxidizable compounds in waste water.

**20. What is meant by blank titration?**

The titration, which is carried out by omitting the sample under consideration, is called blank titration.

**21. What is the purpose of conducting blank titration?**

To find out the amount of potassium dichromate (oxygen) consumed by the organic and inorganic wastes present in water sample.

**22. What is sewage?**

Sewage is commonly a cloudy dilute aqueous solution containing human and household waste water, industrial wastes, ground wastes, street washings. Sewage contains organic and inorganic matters in dissolved, suspended, and colloidal states.

**23. What is meant by industrial sewage?**

The waste water coming out of industrial establishments such as chemical plants, fertilizer

industries, leather tanneries, sugar and paper industries, breweries, textile mills, oil refineries, pharmaceutical units is called an industrial sewage.

**24. What general groups of organic compounds are not oxidized in the COD test?**

Aromatic hydrocarbons and pyridine are not oxidized in COD test.

**25. Explain the colour changes encountered during the titration.**

Ferriin indicator is red in colour in the reduced form with the composition,  $[\text{Fe}(\text{o-Phen})_3]^{2+}$ . When it is added to sewage containing excess of  $\text{K}_2\text{Cr}_2\text{O}_7$ , an oxidizing agent, gets converted into its oxidized form,  $[\text{Fe}(\text{o-Phen})_3]^{3+}$ , which is pale blue (bluish green colour observed). The green colour observed during the course of titration is due to the reduction of  $\text{K}_2\text{Cr}_2\text{O}_7$  by FAS to green  $\text{Cr}_2(\text{SO}_4)_3$ . At the end point, red colour reappears as the indicator is restored to its original form, i.e. reduced form.

**26. What is the limitation of COD?**

One of the chief limitations of COD test is its inability to differentiate between biologically oxidisable and biologically inert organic matter. Also, it does not provide any evidence of the rate at which the biologically active material would be stabilized under conditions that exist in

## Experiment 3

### Estimation of iron in TMT bar by Diphenyl amine indicator method

**AIM:** Determination of the amount of ferrous iron present in the given volume of a solution (100ml) using a standard solution of potassium dichromate.

**PRINCIPLE:**



In the present experiment, ferrous iron to be determined, is oxidized to ferric iron by potassium dichromate in acid medium. The end point is determined by using a redox indicator (for e.g., DPA).

Potassium dichromate is relatively a weak oxidizing agent compared to permanganate and the reaction is slow near the end point. This is mainly due to accumulation of  $\text{Fe}^{+3}$  ions. Syrupy phosphoric acid binds these ions as ferric phosphate and removes them from the reaction sphere, which facilitates the acceleration of the main reaction. The electrode potential of Fe(III)-Fe(II) system is found to be -0.68V in acid medium (0.5M  $\text{H}_2\text{SO}_4$ ) and that of the indicator DPA is -0.76V in 0.5M  $\text{H}_2\text{SO}_4$ . The reduction potential of indicator system is not sufficiently high. Hence the addition of phosphoric acid to lower the reduction potential of Fe(II)--Fe(III) couple by complexation improves end point considerably. As the titration is carried out by addition of  $\text{K}_2\text{Cr}_2\text{O}_7$ , blue-violet colour is obtained at the endpoint.

**Procedure:**

**PART-A: Preparation of 0.05M potassium dichromate solution:**

About 0.735g of  $\text{K}_2\text{Cr}_2\text{O}_7$  is weighed accurately in a clean, dry weighing bottle and transferred into a 250ml volumetric flask through a glass funnel. The substance is dissolved completely in a minimum amount of distilled water and the solution is made up to the mark. The solution is made homogeneous by thorough shaking in the stoppered volumetric flask. The particulars of the weights used are tabulated and the concentration of potassium dichromate is calculated using the given formula.

**PART-B: Determination ferrous in given iron solution**

The given sample of TMT bar solution (ferrous iron solution) is made up to the mark with distilled water (in the given 100 ml volumetric flask) and shaken well to make it homogeneous in concentration. Now, 10 ml of the solution is drawn through a pipette into a 250ml clean conical flask. To this, 20ml of distilled water,

3ml of acid mixture (sulphuric acid and phosphoric acid), 2-3 drops of DPA indicator solution are added with complete mixing. This mixture is then titrated with standard potassium dichromate solution run down from the burette. The contents of the conical flask are swirled throughout the titration till the end point is reached. The end point is determined by a change in colour of the solution from colourless to blue-violet colour. The same procedure of titration is repeated until concurrent readings are obtained and all the observations are tabulated.

Burette: Std. Potassium dichromate solution

Conical flask: 10 ml of TMT bar solution + 20ml of distilled water + 3ml of acid mixture (sulphuric acid and phosphoric acid) + 2-3 drops of DPA

Indicator: 2-3 drops of Diphenyl amine (DPA)

End Point: Blue-violet colour

**Result:** The Amount of ferrous iron [Iron (II)] present in given 100ml solution

= ..... g/ml

### Observation and Calculation:

Preparation of 0.05M potassium dichromate solution:

Weight of weighing bottle +  $K_2Cr_2O_7$  salt =  $W_1$  = .....g

Weight of empty bottle =  $W_2$  = .....g

Amount of  $K_2Cr_2O_7$  salt transferred =  $W_1 - W_2$  = .....g

Normality of  $K_2Cr_2O_7$  solution is=  $\frac{(W_2 - W_1) \times 4}{\text{Equivalent weight of } K_2Cr_2O_7 \text{ (49)}}$

$$= \frac{(W_2 - W_1) \times 4}{49}$$

= .....N1

### Determination ferrous in given iron solution

Burette Reading	I	II	III
-----------------	---	----	-----

Initial B.R			
Final B.R			
Volume of $K_2Cr_2O_7$ solution consumed ( $cm^3$ )			

Volume of  $K_2Cr_2O_7$  solution consumed = .....( $V_1$ )  $cm^3$

$(N_1 \times V_1) K_2Cr_2O_7 = (N_2 \times V_2)$  iron(II) solution

$$N_2 = \frac{N_1 \times V_1}{V_2}$$

$$= \frac{N_1 \times V_1}{10}$$

Amount of ferrous iron (II) present in given 100ml solution =  $\frac{N_2 \times 55.86 \times 100}{1000}$

= ..... g/ml

### **VIVA QUESTIONS & ANSWERS**

**1. What is the aim of the experiment?**

To determine the percentage of iron present in a TMT steel bar using the diphenylamine indicator method.

**2. Principle of the experiment**

Iron in the TMT bar is converted into ferrous ions and oxidized to ferric ions using potassium dichromate. Diphenylamine indicates the end point by a color change.

**3. Why is diphenylamine used as an indicator?**

It is a redox indicator that gives a blue-violet color at the end point.

**4. Which oxidizing agent is commonly used?**

Potassium dichromate ( $K_2Cr_2O_7$ ).

**5. Why potassium dichromate is preferred over potassium permanganate?**

It is a primary standard, stable, available in pure form, and gives a sharp end point.

**6. What is the role of concentrated hydrochloric acid?**

It dissolves iron from the TMT bar and converts it into soluble ferrous chloride.

**7. Why is phosphoric acid added during titration?**

It complexes ferric ions and gives a sharp end point.


**8. What color change is observed at the end point?**

Green to blue-violet.

**9. What type of titration is involved?**

Redox titration.

**10. What is the chemical reaction involved?**



**11. Why is sulfuric acid preferred over nitric acid?**

Nitric acid is an oxidizing agent and may oxidize ferrous ions prematurely.

**12. What is TMT bar?**

Thermo-Mechanically Treated steel bars used in construction.

**13. What percentage of iron is usually present in TMT bars?**

Approximately 98–99%.

**14. Why is the solution heated during dissolution?**

To speed up dissolution of iron.

**15. What precautions should be taken?**

Avoid overheating, use fresh indicator, add indicator near end point, titrate slowly.

**16. What happens if excess indicator is added?**

It may cause early end point and inaccurate results.

**17. Is diphenylamine a self-indicator?**

No, it is an external redox indicator.

**18. Why is the solution cooled before titration?**

High temperature affects indicator behavior and accuracy.

## Experiment 4

### Determination of Total Alkalinity of given water sample

**Theory:** Alkalinity is due to the substances that can cause the formation of hydroxyl ( $\text{OH}^-$ ) ions which in turn can react with strong acids. Alkalinity of a water sample is a measure of its capacity to neutralize acids.

Substances that cause the alkalinity are of three types.

- (i) Hydroxides –  $\text{NaOH}$ ,  $\text{Ca}(\text{OH})_2$ ,  $\text{Mg}(\text{OH})_2$  ;
- (ii) Carbonates –  $\text{Na}_2\text{CO}_3$ ,  $\text{CaCO}_3$ ,  $\text{MgCO}_3$  ;
- (iii) Bicarbonates –  $\text{NaHCO}_3$ ,  $\text{Ca}(\text{HCO}_3)_2$ ,  $\text{Mg}(\text{HCO}_3)_2$ .

Total alkalinity,  $A_t$ , is the sum of the alkalinity due to hydroxides,  $A_h$ , alkalinity due to carbonates,  $A_c$  and the alkalinity due to bicarbonates,  $A_b$ .

$$A_t = A_h + A_c + A_b$$

When a sample of alkaline water is treated with a strong acid such as  $\text{HCl}$ , the following reactions occur.

- (1)  $\text{NaOH} + \text{HCl} \longrightarrow \text{NaCl} + \text{H}_2\text{O}$  (for hydroxides)
- (2)  $\text{Na}_2\text{CO}_3 + \text{HCl} \longrightarrow \text{NaHCO}_3 + \text{NaCl}$  (for carbonates)
- (3)  $\text{NaHCO}_3 + \text{HCl} \longrightarrow \text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$  (for bicarbonates)

Alkalinity is determined by titrating a known volume of water sample against a strong acid using a suitable indicator. Two types of alkalinity are evaluated based on the whether the indicator used is methyl orange or phenolphthalein.

- (i) Alkalinity due to methyl orange,  $A_{m.o}$ . When methyl orange is used, the colour change at the end point is obtained only after all the three reactions given by equations (1), (2) and (3) are complete. Hence methyl orange gives the total alkalinity.

$$A_t = A_{m.o} = A_h + A_c + A_b$$

- (ii) Alkalinity due to phenolphthalein,  $A_{ph}$ . When phenolphthalein is used, the colour change at the end point is obtained after reactions given by equations (1), (2) and (3) are complete and before reaction given by equation (3)

occurs. Thus the alkalinity due to phenolphthalein is due to the hydroxides and half the carbonates as the colour change occurs before the bicarbonates react [see reactions (2) and (3)]. Therefore,

$$A_{ph} = A_h + 0.5A_c$$

### Procedure:

#### Phenolphthalein Alkalinity

- (i) Take 25 ml sample and add two drops of Phenolphthalein indicator, if a slight pink
- (ii) colour appears, it means phenolphthalein alkalinity is present in the water sample.
- (iii) Titrate it with standard HCl (0.02N) slowly and carefully. The pink colour of sample changes and it will become colourless and it is the end point. Record the volume of the titrant consumed as P in ml.

#### Methyl Orange Alkalinity

Test for methyl orange alkalinity is conducted in continuation to the phenolphthalein alkalinity test. In the same sample add two drops of methyl orange indicator. If orange colour develops in the water, it indicates the presence of methyl orange alkalinity. Titrate it with same titrant i.e. HCl (0.02N), the colour of the sample changes from orange to faint pink and that is end point. Record the final reading of burette as T in ml. Find the volume titrant consumed in methyl orange alkalinity by deducting the volume of titrant used in phenolphthalein alkalinity test from the total volume of titrant consumed in both titration. Pipette out 25 ml of water sample into a clean titration flask. Add two drops of methyl orange. Titrate against standard Hydrochloric acid till the colour of the solution changes sharply from yellow to red. Repeat for concordant values

### Observations and Calculations

#### Burette Reading

Trial No.	1	2	3
Final burette reading			
Initial burette reading			
Volume of HCl run down in cm <sup>3</sup>			

The volume of HCl consumed = ..... V cm<sup>3</sup>

Normality of HCl = ..... N

$1000\text{cm}^3$  of 1N HCl = 50g of  $\text{CaCO}_3$  (Equivalent weight of  $\text{CaCO}_3 = 50$ )

$$V\text{cm}^3 \text{ of } X \text{ N HCl is } = \frac{50 \times V \times N}{1000} \text{ g of } \text{CaCO}_3$$

$$= a \text{ g}$$

Total Alkalinity of  $25\text{cm}^3$  of water sample = a g

Total Alkalinity of  $10^6$  (1 million parts)  $\text{cm}^3$  of water sample =  $\frac{a \times 10^6}{25} = z \text{ g of } \text{CaCO}_3$

Therefore total alkalinity of the given water sample = z ppm  $\text{CaCO}_3$  equivalent

**Result:**

Total alkalinity of the water sample = ----- ppm  $\text{CaCO}_3$  equivalent

### VIVA QUESTIONS & ANSWERS

**1. What is alkalinity?**

It is the capacity to neutralize acid. Or it is the capacity of the water to accept the protons.

**2. What are the causes of alkalinity?**

Alkalinity of water is due to presence of  $\text{OH}^-$ ,  $\text{CO}_3^{2-}$ ,  $\text{HCO}_3^-$  ions.

**3. What are the various combinations of anions, responsible for alkalinity**

Carbonates only  $\text{CO}_3^{2-}$

Bicarbonate only ( $\text{HCO}_3^-$ )

Hydroxide only ( $\text{OH}^-$ )

Carbonate and Bicarbonate ( $\text{CO}_3^{2-}$  &  $\text{HCO}_3^-$ )

Carbonate and Hydroxide ( $\text{CO}_3$  &  $\text{OH}^-$ )

**4. Can all the three alkalinity causing ions co-exist? Justify**

Presence of all the three combinations is not possible. Because when  $\text{OH}^-$  &  $\text{HCO}_3^-$  are present in the sample they will react and form  $\text{H}_2\text{O}$  &  $\text{CO}_3^{2-}$  respectively.



Name the types of alkalinity

- i) Phenolphthalein alkalinity
- ii) Methyl Orange alkalinity

**5. What are the indicators used in the determination of alkalinity**

Phenolphthalein and Methyl Orange indicator

**6. What causes Phenolphthalein alkalinity?**

Presence of  $\text{OH}^-$  &  $\text{CO}_3^{2-}$  causes Phenolphthalein alkalinity

**7. Phenolphthalein alkalinity of a water sample is zero. What it indicates?**

It indicates the absence of hydroxyl and carbonate ions in water

**8. How do you express the total alkalinity of water?**

It is expressed in terms of ppm (parts per million) of  $\text{CaCO}_3$

**9. What conclusion you make when phenolphthalein alkalinity is equal to Methyl orange alkalinity?**

When Phenolphthalein alkalinity is equal to methyl orange alkalinity, the alkalinity of water is exclusively due to presence of hydroxyl ions.

**10. What is the working pH of phenolphthalein and methyl orange indicators**

Phenolphthalein =  $\geq 8.3$  (pink)  $\leq 8.3$  (colorless)

Methyl orange =  $\leq 4.3$  (red)  $\geq 4.3$  (yellow)

**11. What is the desirable limit of alkalinity in drinking water according to Indian standards?**

According to Indian standards the desirable limit is up to 200ppm

## Experiment 5

### Determination of acid value of solder flux using KOH

**Aim:** To determine the strength of an acid in a solder flux.

**Principle:** The acid value is the amount of KOH (in mg) required to neutralize the free acid present in

1 gram of solder flux. The free acids in the flux react with standard alkali (KOH or NaOH), and the amount of alkali consumed during neutralization helps to calculate the acid value.

#### Procedure:

##### Part - A: Preparation of Standard NaOH solution:

Weigh accurately 0.8g of the given sodium hydroxide flakes and transfer it into a 100cm<sup>3</sup> standard flask using a funnel. Add distilled water and dissolve the NaOH flakes make it up to the mark and shake well for uniform concentration.

##### Part - B: Estimation of an acid value in solder flux:

Pipette 25 cm<sup>3</sup> of solder flux sample into a clean conical flask. Add about 5 cm<sup>3</sup> of neutralized ethanol or a mixture of ethanol and ether to dissolve the flux. Add 2–3 drops of phenolphthalein indicator to the solution. Titrate the solder flux solution against the standard KOH solution until a persistent pale pink colour appears.

#### Observations and Calculation:

Part – A: Preparation of standard potassium Hydroxide solution:

Weight of the empty weighing bottle + FAS =  $W_1$  = ... g

Weight of the empty weighing bottle =  $W_2$  = ... g

Weight of the FAS =  $W = (W_1 - W_2)$  = ... g

---

Burette Readings	Trail – I	Trail – II	Trail – III
Final Burette Reading			
Initial Burette Reading			
Volume of NaOH solution added in cm <sup>3</sup>			

Volume of KOH solution added in cm<sup>3</sup> (V<sub>2</sub>)=..... cm<sup>3</sup>

**Blank Titration:** Perform a blank titration with the solvent and indicator only to record the volume of alkali used is.. (V<sub>1</sub>) cm<sup>3</sup>

$$\text{Acid Value} = \frac{(V_2 - V_1) \times N_{\text{NaOH solution}} \times \text{Molecular weight of KOH}}{\text{Weight of solder in gram}}$$

Weight of solder in gram

Acid Value =..... KOH/g of solder flux.

**Report:**

---

## VIVA QUESTIONS & ANSWERS

1. What is the acid value of a biofuel?  
The acid value is the amount of potassium hydroxide (in mg) required to neutralize the free fatty acids present in 1 g of biofuel.
2. Why do we determine the acid value of biofuel?  
It indicates the free fatty acid content, which affects fuel quality, stability, and corrosion potential.
3. What is the main reagent used in this experiment?  
0.1 M potassium hydroxide (KOH) solution is used for titration.  
0.2
4. Which indicator is commonly used?  
Phenolphthalein is commonly used as an indicator.
5. What does the pink color signify in the titration?  
It indicates that all free fatty acids have been neutralized and the endpoint is reached.
6. Why is it important to use neutralized alcohol?  
To ensure that the alcohol itself does not react with KOH and affect the titration result.
7. How is the acid value calculated?  
$$\text{Acid value} = (\text{Volume of KOH} \times \text{Normality} \times 56.1) / \text{Weight of biofuel sample (mg KOH/g)}.$$
8. What is an acceptable acid value for biodiesel?  
Typically, less than 0.5 mg KOH/g is considered good quality biodiesel.
9. What happens if the acid value is high?  
High acid value can cause corrosion of engine parts and reduce fuel stability.
10. Can this method be applied to other oils?  
Yes, it can be used for edible oils, fats, and other biodiesel feedstocks to determine free fatty acids.


## Experiment 6

### Estimation Of Acidmixture By Conductometric Sensor


**Aim:** To determine the amount of HCl & CH<sub>3</sub>COOH in the given solution.

**Chemicals:** 0.1 N KCl solution, Standard NaOH solution.

**Theory:** A titration in which the equivalence point or end point of the reaction is determined with the help of measurements of conductivity of the reaction mixture; is known as conductometric titration. The principle underlying conductometric titrations is that the specific conductivity of a solution depends on number and nature of ions, which changes gradually during the titration and lets one to know the end points. Conductivity of unit volume (1 cm<sup>3</sup>) of the electrolytic solution at specified temperature is known as specific conductivity. It is expressed in units of milli Siemens per centimeter (mS cm<sup>-1</sup>). When a mixture of HCl & CH<sub>3</sub>COOH is titrated against NaOH, stronger acid, HCl is neutralized first followed by the neutralization of weak acid, CH<sub>3</sub>COOH. While HCl is neutralized, highly mobile H<sup>+</sup> ions of acid are replaced by less mobile Na<sup>+</sup> ions of the base and specific conductivity decreases. This trend continues until the reach of neutralization of HCl.



After the complete neutralization of strong acid, NaOH added will neutralize weak acid, CH<sub>3</sub>COOH.




Weaker electrolyte, acetic acid is replaced by stronger electrolyte, sodium acetate. Thus, specific conductivity will tend to rise (slowly due to less mobile acetate ion and sodium ion) until the complete neutralization of acetic acid. Later, excessively added NaOH will release relatively more mobile OH<sup>-</sup> ions, which lead to steep rise in specific conductivity.

A plot of specific conductivity against volume of NaOH added will help in knowing the neutralization points. Knowing the normality and volume of NaOH consumed for neutralization, concentration and amount of HCl & CH<sub>3</sub>COOH are calculated.

**Procedure:**

Ensure calibrated condition of the conductivity meter (with the assistance of staff or laboratory instructor). Take Z mL of the given acid mixture in a clean beaker. Dip the conductivity cell in the solution (dilute if required to immerse the conductivity cell properly) and note down the specific conductivity of the solution. Now, add standard NaOH solution from the burette in increments of 0.5 mL. After each addition, stir the solution gently and note down the specific conductivity. As the titration proceeds, the specific conductivity decreases gradually until complete neutralization of HCl and then rises slowly until neutralization of CH<sub>3</sub>COOH. After crossing the neutralization of CH<sub>3</sub>COOH, conductivity rises steeply. Continue the titration for at least 8-10 readings after crossing the neutralization of CH<sub>3</sub>COOH. Plot a graph of specific conductivity against volume of NaOH as shown below. Intersecting lines represent the neutralization points of the two acids. Knowing the normality and amount of NaOH required for neutralization, amounts of HCl & CH<sub>3</sub>COOH are calculated.

**Result:**

Normality of HCl solution = .....N

Amount of HCl = .....g/L

Normality of CH<sub>3</sub>COOH solution = .....N

Amount of CH<sub>3</sub>COOH = .....g/L

**Observations and calculations:**

Solution taken in the burette: Standard NaOH (N NaOH =.....N)

Solution taken in the beaker:

Acid mixture (HCl + CH<sub>3</sub>COOH) = Z = ..... mL + distilled water

for proper immersion of conductivity cell

It implies that  $V_{\text{HCl}} = Z = \dots\dots\dots$  mL;

$V_{\text{CH}_3\text{COOH}} = Z = \dots\dots\dots$  mL

Volume of NaOH (cm <sup>3</sup> )	Specific conductivity, $\kappa$ ( mS cm <sup>-1</sup> )
0.0	
0.5	
1.0	
1.5	
2.0	
2.5	
3.0	
3.5	
4.0	
4.5	
5.0	
5.5	
6.0	
6.5	
7.0	
7.5	
8.0	
8.5	
9.0	
9.5	
10.0	

Volume of NaOH (cm <sup>3</sup> )	Specific conductivity, $\kappa$ ( mS cm <sup>-1</sup> )
10.5	
11.0	
11.5	
12.0	
12.5	
13.0	
13.5	
14.0	
14.5	
15.0	

Volume of NaOH required to neutralize HCl from the graph = V1 =..... mL

Volume of NaOH required to neutralize both HCl & CH<sub>3</sub>COOH from the graph= V2 =..... mL

$$\begin{aligned} \therefore \text{Volume of NaOH required to} & & = V_3 = V_2 - V_1 \\ \text{neutralize only CH}_3\text{COOH} & & = \dots\dots\dots - \dots\dots\dots = \dots\dots\dots \text{ mL} \end{aligned}$$

$$\text{It is known that } N_1V_1 = N_2V_2 \quad \text{or} \quad N_{\text{HCl}} \times V_{\text{HCl}} = N_{\text{NaOH}} \times V_{\text{NaOH}}$$

$$\text{Here, } V_{\text{NaOH}} = V_1$$

$$\begin{aligned} \therefore \text{Normality of HCl} = N_{\text{HCl}} &= \frac{N_{\text{NaOH}} \times V_1}{V_{\text{HCl}}} &= \frac{\dots\dots\dots \times \dots\dots\dots}{\dots\dots\dots} \\ & &= \dots\dots\dots \end{aligned}$$

The amount of HCl in the given solution

$$\begin{aligned} &= N_{\text{HCl}} \times \text{Equivalent weight of HCl (36.5) g/L} \\ &= \dots\dots\dots \times 36.5 \text{ g/L} \\ &= \dots\dots\dots \text{ g/L} \end{aligned}$$

$$\text{Similarly, } N_{\text{CH}_3\text{COOH}} \times V_{\text{CH}_3\text{COOH}} = N_{\text{NaOH}} \times V_{\text{NaOH}}$$

$$\text{Here, } V_{\text{NaOH}} = V_3$$

$$\begin{aligned} \therefore \text{Normality of CH}_3\text{COOH} = N_{\text{CH}_3\text{COOH}} &= \frac{N_{\text{NaOH}} \times V_3}{\text{CH}_3\text{COOH}} \\ &= \frac{\dots\dots\dots \times \dots\dots\dots}{\dots\dots\dots} \\ &= \dots\dots\dots \\ &= N_{\text{CH}_3\text{COOH}} \times \text{Equivalent weight of CH}_3\text{COOH (60) g/L} \end{aligned}$$

$$= \text{-----} \times 60 \text{ g/L}$$

The amount of  $\text{CH}_3\text{COOH}$  in the given solution = .....g/L

### **VIVA QUESTIONS and ANSWER**

**1. What is conductance? Mention its unit.**

The reciprocal of resistance is called the conductance. The unit is  $\text{ohm}^{-1}$  or mho or Siemens (S).

**2. Mention the factors on which the conductance of solution depends.**

The conductance of solution depends on

- Number of ions
- Charge on the ions
- Mobility of ions
- Temperature

**3. How does conductance of solution vary with temperature?**

It increases with increase of temperature (unlike in the case of metallic conductors).

**4. Name the current carriers in the case of electrolytes.**

The ions present in the solution.

**5. What is specific conductivity?**

It is the conductivity of the solution placed between two electrodes of  $1\text{cm}^2$  area of cross section, which are  $1\text{cm}$  apart. It is expressed in  $\text{ohm}^{-1}\text{cm}^{-1}$ .

**6. Define cell constant.**

It is the ratio of distance between the two electrodes to the area of cross section.

$$\text{Cell Constant} = l/a \qquad \text{Its unit is } \text{m}^{-1}.$$

**7. What is meant by Conductometric titration?**

The estimation of substances in solution by observing sudden change in conductance of the solution at equivalence point is known as Conductometric titration.

**8. Mention the types of conductivity measurements.**

Specific conductivity, molar conductivity and equivalent conductivity.

**9. What is meant by molar conductivity?**

It is the conductance produced by all the ions present in one litre of solution containing gram molecular weight of the electrolyte.

**10. What is meant by equivalent conductivity?**

It is the conductance produced by all the ions present in one litre of solution containing gram equivalent weight of the electrolyte.

**11. What is conductivity cell?**

It is a device used to measure the conductance of solution. It consists of two platinum foils

**12. Account for the followings:****Conductance of the solution decreases in the beginning of titration.**

It is due to the replacement of high mobile  $H^+$  ions of HCl by less mobile  $Na^+$  ions of NaOH solution.

**Conductance of the solution increases slowly in the middle of the titration (i.e. after the first equivalence point).**

It is due to the neutralization of acetic acid by NaOH, which leads to the formation of sodium acetate salt, which ionizes further to give  $CH_3COO^-$  and  $Na^+$  ions.

**Conductance of the solution increases after the end point.**

It is due to the presence of excess of  $OH^-$  ions of NaOH.

**The slope of the line in the beginning of titration is higher than that of the line after the end point**

It is due to the high mobility of  $H^+$  ions HCl (which are replaced by less mobile  $Na^+$ ) when compared to the low mobile  $OH^-$  ions of NaOH.

**13. What are the advantages of Conductometric titrations?**

- i) The method is accurate in dilute as well as more concentrated solutions.
- ii) It can also be employed with colored solutions.
- iii) Mixture of acids can be titrated more accurately.

Ohm's law states that the current,  $I$  (ampere), flowing in a conductor is directly proportional to the applied electromotive force,  $E$  (volt) and inversely proportional to the resistance,  $R$

**14. State ohm's law.**

(ohm) of the conductor.

$$I = E/R$$

**15. What is conductivity?**

The reciprocal of resistivity is called conductivity.

**16. Which of the above conductance measured during conductometric titration?**

---

The specific conductance is measured.

**17. What is a cell?**

A device which produces an electromotive force and delivers an electric current as the result of a chemical reaction is known as a cell.

**18. What is the principle involved in conductometric titration?**

In conductometric titration, there is a sudden change in conductance of solution near the end point. Hence the end point is determined graphically by plotting conductance against titre values. The principle underlying conductometric titration is the replacement of ions of a particular conductance by ions of different conductance during the titration.

**19. How is the equivalence point obtained in conductometric titration?**

During the progress of the titration, changes in conductivity occur. The conductivity is measured after each addition of a small volume of the titrant. A graph of conductivity (on Y-axis) versus volume of titrant (on X-axis) is plotted when two or more straight lines are obtained. The point of intersection of the two straight lines gives the equivalence point.

**20. In the titration of a mixture of acids (HCl and CH<sub>3</sub>COOH) with a strong base (NaOH), the conductance first decreases, then rises steadily and finally rises steeply. Why?**

Upon adding a strong base to a mixture of a strong acid and a weak acid, the conductance falls due to the replacement of highly mobile H<sup>+</sup> ions of the strong acid (HCL) by less mobile Na<sup>+</sup> ions of the base. The conductance falls till all the H<sup>+</sup> ions are replaced (i.e. till HCL is neutralised completely). The conductance then rises steadily as the weak acid is converted into its salt. Finally, the conductance rises steeply as excess of alkali is introduced


## Experiment 7

### Estimation of Fe in Rust sample by Potentiometric sensor

**Aim:** To determine the amount of FAS in the given solution


**Chemicals:** Dilute  $\text{H}_2\text{SO}_4$ , Standard  $\text{K}_2\text{Cr}_2\text{O}_7$ , FAS

**Theory:** A titration in which the equivalence point of the reaction is determined with the help of measurements of cell potentials is known as potentiometric titration. A suitable cell made of indicator electrode and reference electrode is constructed in known volume of the test solution. For the titration of FAS against  $\text{K}_2\text{Cr}_2\text{O}_7$ , platinum electrode is used as indicator electrode and calomel electrode as reference electrode. Indicator electrode (Pt) potential is a function of the redox system and activities (or concentration) of the respective ionic species in the solution. Reference electrode (calomel) potential remains constant during titration. For a redox system, same metal with two oxidation states, will compete for mutual conversions,




$$E = E^0 + 0.0591/n \log [(\text{Oxidized species}) / (\text{Reduced species})]$$

For the titration of FAS against  $\text{K}_2\text{Cr}_2\text{O}_7$ , Ferrous ion undergoes oxidation to ferric state:



Dichromate ion undergoes reduction to chromic state:




Platinum electrode potential is decided by the redox system. Before the end point, the electrode is  $\text{Pt} | (\text{Fe}^{3+}, \text{Fe}^{2+})$ . During the titration,  $\text{Fe}^{3+}$  ion concentration increases and that of  $\text{Fe}^{2+}$  ion decreases until the reach of equivalence point. The numerical value of the ratio,  $[\text{Fe}^{3+}]/[\text{Fe}^{2+}]$  increases and potential increases After the end point, the electrode changes as  $\text{Pt} | (\text{Cr}_2\text{O}_7^{2-}, \text{Cr}^{3+})$  resulting in abrupt change in electrode potential and consequently cell-potential at the end point. The cell-potential is decided by the ratio  $[\text{Cr}_2\text{O}_7^{2-}] / [\text{Cr}^{3+}]$  Sudden change in cell potential helps in the determination of equivalence point. With the

knowledge of normality and volume of  $K_2Cr_2O_7$  consumed the amount of FAS is calculated.

**Procedure:**

Standardize the potentiometer using electronic standard cell on the instrument (potential needs to be set to 1.018 V or 1018 mV using the calibration knob). Take suggested volume, Z mL of FAS solution from a burette into a clean beaker. Add 10 mL of dilute sulfuric acid. Wash the platinum and saturated calomel electrode assembly and place it in the solution. Add requisite amount of distilled water required for immersion of cell assembly. Measure the EMF of the experimental cell on the potentiometer. Add  $K_2Cr_2O_7$  solution from a burette in increments of 0.5 mL and stir the solution thoroughly. Measure the EMF after each addition. Equivalence point is recognized by sudden jump in EMF. Continue the addition of  $K_2Cr_2O_7$  until 6 – 8 readings beyond the equivalence point. Plot a graph of  $\Delta E/\Delta V$  against the volume of  $K_2Cr_2O_7$  (differential curve is obtained). Determine the equivalence point from the graph as shown in figure below. Calculate the strength and amount of FAS.



**Result:**

The normality of FAS solution = .....N;

The amount of FAS in the solution = .....g/L



## VIVA QUESTIONS & ANSWERS

### 1. What is a Potentiometric titration?

Ans. The determination of equivalence point of red-ox titration on the basis of potential measurements is known as Potentiometric titration.

### 2. What are the electrodes used in Potentiometric titration?

Ans. Platinum and calomel electrodes.

### 3. What is the principle Potentiometric titration?

Ans. The principle is the measurement of the emf between the platinum electrode (indicator electrode) and a reference electrode (calomel electrode). In this titration, the measurements of potential are made while the titration is in progress. The equivalence point is sudden change in the potential.

### 4. What is an indicator electrode?

The electrode, whose potential is dependent on the concentration of the ion being determined,.


### 5. What is the determining factor in the red-ox titration?

Ans. The determining factor is the ratio of concentrations of oxidized species to reduced species.

### 6. What is the reaction-taking place between FAS and Dichromate?

Ans. Acidified potassium dichromate oxidizes ferrous to ferric.

### 7. Give the reaction between FAS and K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> in Potentiometric titration. Acidified K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> oxidizes ferrous sulphate to ferric sulphate and itself gets reduced to chromic sulphate.



### 8. Mention the advantages of Potentiometric titrations.

1. Turbid, colored solutions can be titrated.
2. Mixture of solutions or very dilute solutions can be titrated.
3. The results are more accurate since the end point is determined graphically.

### 9. What is the purpose of adding dilute sulphuric acid?

This is because potassium dichromate acts as oxidizing agent only in acidic media.

### 10. What is single electrode potential?

The potential that is developed when an element is in contact with a solution containing its own ions is called single electrode potential

### 11. What is standard electrode potential?

The potential that is developed when an element is in contact with a solution containing its own ions of 1 M concentration at 298 K is referred to as standard electrode potential. If the gases are involved, they must be passed at a partial pressure of 1 atmosphere.

### 12. What is meant by e.m.f.?

E.M.F. is the potential difference required to drive a current across the electrodes.

E.M.F. =  $E_{\text{cathode}} - E_{\text{anode}}$

### 13. What is a potentiometer?

It is a device or circuit used for comparing potential sources.

### 14. Why is the beaker solution gradually changes into green during the course of titration? When

FAS in beaker reacts with K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>, it reduces K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> to Cr<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> which is a green salt

solution.

**15. Why is the EMF rises steeply soon after the equivalence point?**

This is because, the potential of the solution before the equivalence point is determined by  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$  system only i.e., 0.75V, while at equivalence point, it is determined by both  $\text{Fe}^{3+}$  and  $\text{Cr}_2\text{O}_7^{2-}$  ions which is = 1.04V. But beyond equivalence point, the potential of the solution is determined by  $\text{Cr}_2\text{O}_7^{2-}/\text{Cr}^{3+}$  only i.e. = 1.33V. Therefore, just after the equivalence point, the potential of the solution rises steeply.

## Experiment 8


### Determination of pK<sub>a</sub> of vinegar using pH sensor (Glass electrode)

**Aim:** To determine the pK<sub>a</sub> value of the given weak acid

**Chemicals:** Standard buffer solutions (pH = 7.0, 4.0, 9.2), NaOH solution

**Theory:** pK<sub>a</sub> is a measure of the strength of the acid. Higher the pK<sub>a</sub>, weaker is the acid. It is defined as negative logarithm of dissociation constant, K<sub>a</sub> [pK<sub>a</sub> = -logK<sub>a</sub>]. pK<sub>a</sub> is determined by the measurement of variation in pH during titration of the acid against a base.


For the titration of weak acid against strong base, the change in pH is lesser until the reach of equivalence point. For example, pH changes slowly until neutralization point during the titration of CH<sub>3</sub>COOH against NaOH. Reaction that takes place during the titration is



During the titration, solution will become a mixture of weak acid (unreacted CH<sub>3</sub>COOH) and its salt (CH<sub>3</sub>COONa), which represents the buffer. Therefore, pH changes slowly until the neutralization. At the equivalence point, the pH rises rapidly owing to removal of H<sup>+</sup> ions (acid) and addition of OH<sup>-</sup> ions (alkali) [the buffer disappears at this point of titration]. The change in pH during the titration is described by Henderson-Hasselbatch equation.

$$\text{pH} = \text{pK}_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$

At half equivalence point, half the acid will have reacted to give salt; and half the acid will have remained unreacted. Therefore, [Salt] = [Acid] and log(1) = 0. Thus, above equation becomes, pH = pK<sub>a</sub> i.e., pH at half equivalence point gives pK<sub>a</sub> value of the weak acid. pH is measured on a pH meter using combined glass electrode (cell constructed by the coupling of glass electrode & Ag-AgCl electrode). The cell used is represented as



The cell potential evaluated is a measure of H<sup>+</sup> ion concentration in the solution, which is read in terms of pH. The graphs ΔpH/ ΔV vs Volume of NaOH (V<sub>NaOH</sub>) and pH vs V<sub>NaOH</sub> together will assist in the evaluation of pK<sub>a</sub>.

#### Procedure:

Ensure calibrated condition of the conductivity meter (with the assistance of staff or laboratory instructor). Take the suggested volume (Z mL) of weak acid into a clean beaker. Place the combined glass electrode in the weak acid. Add sufficient quantity of distilled water to ensure proper immersion of electrode in the solution and record the pH value. Add NaOH solution from the burette in increments of 0.5 mL every time and stir the solution well. Note down the pH after

each addition. Equivalence point is recognized by sudden jump in pH. Continue the addition of NaOH until 6 – 8 readings beyond the equivalence point. Plot a graph of  $\Delta\text{pH}/\Delta V$  vs Volume of NaOH ( $V_{\text{NaOH}}$ ) find out the equivalence point (Fig.1). Plot another graph of pH versus volume of NaOH, mark the position of half equivalence point and obtain  $\text{pK}_a$  value (Fig.2).




Fig.1




Fig.2

**Result:**

$\text{pK}_a$  value of the given weak acid = .....



Ans.  $pK_a = -\log_{10}K_a$ , where  $K_a$  is dissociation constant of weak acid

**3. What is meant by pH of a solution?**

Ans.  $pH = -\log_{10} (H^+)$

**4. What are the electrodes used in pH meter?**

Ans. Glass electrode and calomel electrode.

**5. Why glass electrode is known as ion selective electrode?**

Ans. This is because it is able to respond to  $H^+$  ions only and develop potential while ignoring the other ions present in the solution.

**6. How is the measurement of pH made?**

Ans. Determining the EMF of the cell containing glass electrode and calomel electrode immersed in the

solution makes the measurement of pH. The emf of the cell is expressed as  $E = K + 0.0591pH$ .

**7. How pH and pKa are related?**

Ans.  $pH = pK_a + \log_{10} (\text{salt}) / (\text{acid})$

The above equation is known as Henderson-Hasselbatch equation. At half Equivalence point (salt) = (acid) and therefore  $pH = pK_a$

**8. Why does pH increase suddenly after the equivalence point?**

Ans. At the equivalence point, the base neutralizes the weak acid. Afterwards, the concentration of hydroxyl ions increases, resulting in sudden increase of pH.

**9. Why is (acid) = (salt) at half equivalence point?**

Ans. At half equivalence point, half of the acid is converted into its salt.

**10. How are pKa and strength of weak acid related?**

Ans. Higher the  $pK_a$ , lower will be the strength of the acid and vice versa.

**11. What is the dissociation constant of weak acid?**

Ans. The dissociation constant of weak acid  $K_a = (H^+) (A^-) / (HA)$

## Experiment 9

### Estimation Of Copper Present In E-Waste By Optical Sensor

**Aim:** To determine the amount of copper in the given test solution

**Chemicals:** Standard solution of CuSO<sub>4</sub> (0.05 N), 1:1 Ammonia solution

**Theory:** An analysis by the way of measurements of absorbance of light in the visible range is colorimetry. It depends upon the measurement of quantity of light absorbed by a coloured solution. The wavelength of light absorbed is a function of nature (quality) of the substance and the amount of light absorbed is a function of concentration (quantity).

When certain light with intensity  $I_0$  is passed through a transparent solution, part of it is absorbed ( $I_a$ ), a part is reflected ( $I_r$ ) and another part is transmitted ( $I_t$ );  $I_0 = I_a + I_r + I_t$

For an interface of air and glass, the amount of light reflected is negligible;

$$I_0 = I_a + I_t$$

Quantitative measurements of absorption are based on Beer's law and Lambert's law or Beer-Lambert's law. **Beer's law** states that „when a monochromatic beam of light (light of single wavelength) passes through a transparent medium, the intensity of the emitted light decreases exponentially as the concentration of the medium increases arithmetically“. **Lambert's law** states that „when a monochromatic beam of light passes through a transparent medium, the intensity of the emitted light decreases exponentially as the thickness of the medium increases arithmetically“. Combined **Beer-Lambert's law** infers that, „when a monochromatic beam of light passes through a transparent medium, the amount of light absorbed by coloured solution is proportional to the concentration and thickness of the light absorbing medium“. It can be expressed as  $A = \log [I_0 / I_t] = \epsilon c t$  where, „ $I_0$ “ is intensity of incident light, „ $I_t$ “ is intensity of transmitted light, „ $t$ “ is thickness of the light absorbing medium (cm), „ $c$ “ is concentration of the coloured constituent in the solution (mol/L) and „ $\epsilon$ “ is molar extinction coefficient (amount of light absorbed when „ $t$ “ and „ $c$ “ are 1 cm and 1 mol/L respectively). Since,  $\epsilon$  is a constant, absorbance is proportional to the concentration of coloured constituent in solution, provided the thickness of the light absorbing medium is kept constant. Mathematically,  $A \propto c$

Reliability of colorimetric analysis requires appreciable colour intensity of solution. Therefore, in the estimation of copper, light blue coloured Cu<sup>2+</sup> is reacted with ammonia to give intense blue coloured tetraamine copper(II) complex.



**Observations and calculations:**

Normality of  $\text{CuSO}_4$  solution (given) = 0.05N

Strength of Ammonia solution used = 1:1

Sl. No.	Volume of $\text{CuSO}_4$ (V mL)	Volume of 1:1 $\text{NH}_3$ (mL)	Concentration of Copper $N_S = \frac{V \times 0.05}{50}$ $= V \times 10^{-3} \text{ N}$	Absorbance (or O.D.)
1	0 (Blank solution)	4	0	0
2	2	4	$2 \times 10^{-3}$	
3	4	4	$4 \times 10^{-3}$	
4	6	4	$6 \times 10^{-3}$	
5	8	4	$8 \times 10^{-3}$	
6	10	4	$10 \times 10^{-3}$	
7	Test solution	----	----	

Concentration of copper in the test solution from the graph

$$= N_{\text{Cu}} = \dots\dots\dots \times 10^{-3} \text{ N}$$

Amount of copper in the test solution

$$= N_{\text{Cu}} \times \text{Equivalent weight of copper (63.54) g/L}$$

$$= N_{\text{Cu}} \times 63.54 \text{ g/L}$$

$$= \dots\dots\dots \times 63.54 \text{ g/L}$$

$$= \dots\dots\dots \text{g/L}$$

### **VIVA QUESTIONS & ANSWERS**

**1. What is colorimetry?**

Ans. It is Chemical analysis through measurements of absorption of light radiation in the visible region of spectrum (400-700nm) with respect to known concentration of the substance.

**2. What forms the basis for colorimetric determination?**

Ans. The variation in the absorption of light with change in concentration of the substance is the basis of calorimetric determination.

**3. What is photoelectric colorimeter?**

Ans. It is an electric instrument, which measures the amount of light absorbed using a photocell.

**4. What are filters? Why are they used?**

Ans. The filters consist of thin films of gelatin containing different dyes or different colored glasses. This is used in colorimeters to select any desired spectral region. (Wavelength)

**5. What is frequency?**

Ans. It is the number of waves passing through a point per second. It is represented as  $\nu$ .

**6. What is wave number?**

Ans. It is reciprocal of wavelength.  $1/\lambda$   
represents it  $1/\lambda = \nu / c = \text{Frequency} / \text{velocity of light}$

**7. State Beer's law.**

Ans. The intensity of transmitted light decreases exponentially as the concentration of the absorbing substance increases arithmetically.

**8. State Lambert's law.**

Ans. The intensity of transmitted light decreases exponentially as the thickness of the absorbing medium increases arithmetically.

**9. State Beer-Lambert law.**

Ans. The amount of light absorbed is proportional to concentration and thickness of the medium.  $A = \log I_0 / I_t = \epsilon ct$

$\epsilon$  is Molar extinction coefficient;  $c$  is concentration;  $t$  is path length

**10. Why different volumes of solutions are taken?**

Ans. Different volumes are taken to get calibration graph.

**11. What is blank solution?**

Ans. Solution without the test solute is known as blank solution.

**12. Why ammonia is added to copper solution? Why is that the same amount of ammonia added?**

Ans. Ammonia is added to get cuproammonium sulfate, a dark blue complex. Same amount of ammonia is added to nullify the absorbance due to any coloring impurities present in ammonia.

**13. Why is the estimation of copper done at 620 nm wavelengths?**

Ans. It is done because; the complex shows a maximum absorbance at 620 nm. ( $\lambda_{\text{Max}}$ )

**14. What is meant by transmittance?**

Ans. It is the ratio of intensity of transmitted light ( $I_t$ ) to that of the incident light

( $I_0$ )  $T = I_t / I_0$

**15. What is absorbance or optical density?**

Ans. It is reciprocal transmittance.  $A = 1 / T = I_0 / I_t$

**16. How optical density is related to the concentration of the substance?**

Ans The optical density is directly proportional to concentration of the substance.

**17. Mention a few important criteria for satisfactory colorimetric analysis**

- I. The solute should not undergo salvation, association, hydrolysis or polymerization in the solvent used.
- II. The color produced should be sufficiently stable
- III. Clear solutions free from traces of precipitates or foreign substances in either blank or standard solutions

**18. Mention a few advantages of colorimetric determinations.**

Ans. Colorimetric method will often give more accurate results at low concentrations. It is more useful for biological substances.

**19. How is optical density related to the concentration of the substance?**

The optical density is directly proportional to the concentration of the substance.

## Experiment 10

### Determination of Viscosity coefficient of Coolant Samples

**Aim:** To determine the viscosity coefficient of the given organic liquid.

**Chemicals:** Distilled water

**Theory:** A liquid flowing through cylindrical tube of uniform diameter is expected to move in the forms of molecular layers. A layer close to the surface is almost stationary, while that at the axis of the tube moves faster than any other intermediate layer. A slow moving layer exerts a drag or friction on to the nearest layer. This property that retards or opposes the motion is called viscosity.

The viscosity coefficient of viscosity is defined as the tangential force per unit area required to maintain a unit velocity gradient between any two successive layers of a liquid situated a unit distance apart.

The viscosity coefficient of a liquid is given by Poiscuille's formula,

$\eta = \frac{\pi r^4 \rho}{8lv}$  where,  $v$  is volume of the liquid,  $r$  is the radius of the tube,  $l$  is length of the tube,  $\rho$  is the pressure difference between two ends of the tube,  $\eta$  is the coefficient of viscosity of the liquid.

If equal volumes of liquids are allowed to flow through the same tube under identical conditions, then,

$$\eta / \eta_w = t_l \times d_l / t_w \times d_w \quad (1)$$

The time taken ( $t_l$ ) by the test liquid to flow through the capillary is determined. The time taken ( $t_w$ ) by water to flow through the same capillary is measured under identical conditions. Thus, the viscosity coefficient of test liquid is calculated from the densities ( $d_l$ ,  $d_w$ ) of the test liquid and water and the viscosity coefficient ( $\eta_w$ ) of water. Viscosity is temperature dependent and hence the experiment is carried out at constant temperature.

#### Procedure:

Fix an Ostwald's viscometer vertically in a thermostat (water contained in a beaker; note down the temperature of water). Take appropriate volume (say, 25 mL) of the given liquid into wider limb of viscometer. Draw it up in the capillary limb slightly higher than the upper mark above the glass bulb

Allow the liquid to flow down. Start a stop-watch when the lower meniscus of liquid crosses the upper mark (above the bulb) and stop it when the lower meniscus crosses the lower mark (below the bulb).

Read & record the flow time in seconds. Repeat the measurements for concordant values. Wash the viscometer, rinse with acetone and dry it in a hot air oven. Cool the viscometer thoroughly and fix it vertically in the thermostat. Take identical volume (25 mL) of distilled water in the viscometer and measure its flow time. Knowing the density & viscosity of water, density of liquid and flow times of water & liquid under experimental temperature, viscosity of the liquid is calculated.

**Result:** The viscosity coefficient of the conductive ink is----- milli poise

**Observations and calculations:**

Temperature of the thermostat = °C

Density of test liquid (given)  $d_l = \dots\dots\dots \text{g/cm}^3$

Density of distilled water (given)  $d_w = \dots\dots\dots \text{g/cm}^3$

Viscosity of distilled water (given)  $\eta_w = \dots\dots\dots \text{mP}$

**Flow time measurement:**

T	Flow time (seconds)			
	Trial - 1	Trial - 2	Trial - 3	Average (S)
Test liquid				$t_l =$
Distilled Water				$t_w =$

flow of test liquid  $t_l = \dots\dots\dots \text{S}$

Time of flow of water  $t_w = \dots\dots\dots \text{S}$

Viscosity coefficient of the test liquid =  $\eta_l = \frac{(\eta_w \times d_l \times t_l)}{(d_w \times t_w)} \text{ mp}$

## **VIVA QUESTIONS & ANSWERS**

**1. What is viscosity?**

The internal friction between the moving layers of molecules of liquid is called viscosity i.e. a slow moving molecular layer will exert a backward pull on the fast moving molecular layer.

**2. Define density of liquid. Mention its unit.**

It is defined as the ratio of mass per unit volume of the liquid.

$$\text{Density} = \text{Mass/Volume}$$

It is expressed in terms of Kg/m<sup>3</sup>

**3. Define viscosity coefficient of liquid.**

It is defined as tangential force per unit area required to maintain a unit velocity gradient between any two successive layers of liquid.

**4. Mention the CGS and SI units of viscosity coefficient.**

CGS unit – milli poise SI

unit – Nsm<sup>-2</sup>

**5. How do you convert viscosity coefficient in CGS unit into SI unit?**

$$1 \text{ milli poise} = 10^{-4} \text{ Nsm}^{-2}.$$

**6. Give the Poiseuille's equation.**

$$\eta = \frac{\pi Pr^4 t}{8 V l}$$

P = Hydrostatic pressure; r = radius of capillary tube; t = time for flow of liquid; V = volume of liquid and l = length of the tube.

**7. How does viscosity vary with temperature?**

It decreases with increase of temperature.

**8. Why should Viscometer be dried before the time flow measurements of the liquid?**

To avoid the formation of emulsion, which changes the time for flow of liquid.

**9. How do you dry the Viscometer?**

It is dried first by rinsing with acetone and then keeping in oven.

**10. Why acetone is used for cleaning Viscometer?**

Since acetone is a volatile liquid, it is used to dry the Viscometer quickly.

**11. Why is the viscosity measurements carried out at lab temperature?**

Physical constants like density and viscosity of liquid vary with temperature.

**12. Mention the factors, which affect the viscosity of the liquid.**

1. Viscosity increases with increase in molecular weight.
2. Viscosity decreases with increase of temperature.

3. Polar compounds are more viscous than non polar compounds.

**13. The bulk of the viscometer is immersed in thermostat (beaker filled with water) during the course of the experiment. Why?**

To maintain a constant temperature, since the viscosity vary with change in temperature.

**14. What is density of a liquid?**

The density of a liquid is its mass divided by its volume.

**15. The density of a substance is expressed relative to what?**

The density of a substance is expressed relative to that of water at 4° C.

**16. What is specific gravity?**

Specific gravity or the relative density is the weight of a given liquid divided by the weight of an equal volume of water at the same temperature

## Experiment 11

### Interpretation of pKa values of a cooling fluid using origin software.

**Aim:** To interpret experimentally obtained titration data of a weak acid and determine its pKa value accurately using graphical and mathematical analysis tools available in Origin.

**Principle:** The interpretation of pKa from titration data is based on the characteristic behavior of a weak acid– strong base titration curve. When pH is plotted against the volume of base added, a sigmoidal curve is obtained. The curve contains important regions: the buffer region, the half-equivalence point, and the equivalence point.

According to the Henderson–Hasselbalch equation:

$$pH = pKa + \log \left[ \frac{[Salt]}{[Acid]} \right]$$

At half equivalence point, [salt] = [acid] and hence pH = pKa. Thus pH at half equivalence point gives the pKa of weak acid.


Thus, correct interpretation of the titration curve allows identification of:

- Equivalence point (maximum slope)
- Half-equivalence point (half of equivalence volume)
- Corresponding pH at half-equivalence, which equals pKa

Origin software enables precise graphical visualization, derivative analysis, and curve fitting, which improves accuracy compared to manual estimation.

### Procedure


First, the experimental titration data (volume of NaOH vs. pH) is entered into two columns in Origin. The first column is designated as the X-axis (volume of base added), and the second column as the Y-axis (pH values).




The screenshot shows the Origin software interface with a data table. The table has two columns: 'Volume of NaOH' (X-axis) and 'pH' (Y-axis). The data points are as follows:

Volume of NaOH	pH
2	2.06
3	2.1
4	2.2
5	2.35
6	2.43
7	2.61
8	2.94
9	3.34
10	3.45
11	3.66
12	3.87
13	4.85
14	4.89
15	8.47
16	10.01
17	10.53
18	11
19	11.1
20	11.13
21	11.15

Next, a titration curve is generated by selecting both columns and choosing a 2D line plot (pH vs. Volume). The resulting graph is carefully examined to observe the buffer region and the steep vertical rise near the equivalence point.



To determine the equivalence point accurately, the first derivative of the data ( $\Delta\text{pH}/\Delta V$ ) is calculated using the “Differentiate” tool under the Analysis menu. A new plot of the first derivative versus volume is generated. The peak of this derivative curve corresponds to the equivalence volume. Once the equivalence volume is identified, the half-equivalence volume is calculated by dividing the equivalence volume by two. Using the original titration curve, the pH corresponding to this half-equivalence volume is determined either by direct graph reading or by using the “Interpolate/Find X-Y” tool in Origin for precise value extraction.



This pH value at half-equivalence is recorded as the pKa of the weak acid.

**Result:** The interpreted pKa value is compared with literature values to evaluate accuracy and reliability of the experimental data.

## Experiment - 12

### Chemical structure drawing using software: ChemDraw/ChemSketch

**Aim:** Chemical Structure Drawing Using ChemDraw


#### Procedure

Chem Draw is a widely used chemical drawing program designed for creating accurate molecular structures, reaction mechanisms, laboratory diagrams, and publication-quality chemical illustrations. It is extensively used in academic laboratories, pharmaceutical research, and chemical industries because it ensures structural accuracy, correct bond angles, proper valency representation, and standardized formatting.

#### Important Tools in ChemDraw

ChemDraw provides several essential tools that facilitate chemical drawing:

- **Bond Tools** – Used to draw single, double, triple, and wedge/dash bonds.
- **Ring Tool** – Inserts common ring systems such as benzene, cyclohexane, and other cyclic structures.
- **Atom/Text Tool** – Allows insertion or modification of atoms (O, N, Cl, etc.).
- **Selection Tool (Arrow Tool)** – Used to move, rotate, resize, or select structures.
- **Reaction Arrow Tool** – Used for drawing reaction schemes.
- **Clean Up Structure Option** – Automatically adjusts bond angles and geometry to standard values.
- **Generate Name Option** – Provides the IUPAC name of the drawn structure.



## Example: Drawing the Structure of Aspirin ( $C_9H_8O_4$ )

Aspirin, chemically known as acetylsalicylic acid, contains:

- A benzene ring
- A carboxylic acid group ( $-COOH$ )
- An ester functional group ( $-OCOCH_3$ )

### Step-by-Step Drawing Procedure


First, ChemDraw is opened and a new document is created using **File** → **New**. The benzene ring is drawn by selecting the **Ring Tool** and choosing the benzene template. Clicking on the drawing canvas inserts a six-membered aromatic ring with alternating double bonds.

Next, the **carboxylic acid group** ( $-COOH$ ) is attached to one carbon of the benzene ring. A single bond is drawn outward from the ring carbon. At the end of this bond, a carbon atom is present. Using the **Double Bond Tool**, a double bond is added to an oxygen atom ( $=O$ ). Then, another oxygen is attached with a single bond, and a hydrogen atom is added to this oxygen to complete the  $-COOH$  group.

After this, the **ester group** ( $-OCOCH_3$ ) is added to the adjacent (ortho) carbon of the benzene ring. A single bond is drawn from the ring carbon and an oxygen atom is inserted. From this oxygen, a carbonyl carbon is added, followed by a double-bonded oxygen ( $=O$ ). Finally, a methyl group ( $-CH_3$ ) is attached to the carbonyl carbon using the single bond tool.

Once the structure is complete, the entire molecule is selected using the **Selection Tool**, and the **Structure** → **Clean Up Structure** option is applied. This automatically adjusts bond angles and spacing to standard geometry, producing a neat and publication-ready structure.

The completed structure is then saved using **File** → **Save As** in the required format such as *.cdx*, *.pdf*, or *.png*.



**Result:** The structure of aspirin (acetylsalicylic acid) was successfully constructed using ChemDraw, demonstrating correct use of bond tools, ring tools, functional group insertion, and structure cleanup features.

# Lab Manual 2

## PYTHON PROGRAMMING

### (1BPLC105B/205B)

#### CO's And PO's Mapping Chart

Sl. No.	Description	PO 1	PO 2	PO 3	PO 4	PO 5	PO 6	PO 7	PO 8	PO 9	PO 10	PO 11	PSO 1	PSO 2
1	CO1: Develop scripts using primitive language constructs of python.	3	2			3							1	
2	CO2: Identify the methods to manipulate primitive python data structures	3	1			2							1	
3	CO3: Make use of Python standard libraries for programming.	3	2			3							2	
4	CO4: Build scripts for performing file operations.	3	2			3							1	
5	CO5: Illustrate the concepts of Object-Oriented Programming as used in Python.	3	1			2							1	

#### Evaluation:

<b>Integrated Lab(IPCC)</b>		
<b>CIE</b>		
<b>Particulars</b>	<b>Marks</b>	<b>Total</b>
Performance	07	15
Journal	05	
Viva Voce	03	
<b>Lab IA</b>		
<b>Particulars</b>	<b>Marks</b>	<b>Total</b>
IA	100	10 (Reduced)
Total (CIE +Lab IA)		25

**Mapping of Experiments with CO, PO and PSO**

S.No	Experiment Details	CO	PO	PSO
1a	Develop a python program to read 2 numbers from the keyboard and perform the basic arithmetic operations based on the choice. (1-Add, 2-Subtract, 3-Multiply, 4-Divide).	CO1	PO1	PSO1
1b	Develop a program to read the name and year of birth of a person. Display whether the person is a senior citizen or not.	CO1	PO2	PSO1
2a	Develop a program to generate Fibonacci sequence of length (N). Read N from the console.	CO1	PO1	PSO1
2b	Write a python program to create a list and perform the following operations: Inserting an element, Removing an element, Appending an element, Displaying the length of the list, Popping an element, Clearing the list.	CO2	PO1	PSO1
3a	Read N numbers from the console and create a list. Develop a program to print mean, variance and standard deviation with suitable messages.	CO2	PO2	PSO1
3b	Read a multi-digit number (as chars) from the console. Develop a program to print the frequency of each digit with a suitable message.	CO2	PO2	PSO1
4	Develop a program to print 10 most frequently appearing words in a text file. [Hint: Use a dictionary with distinct words and their frequency of occurrences. Sort the dictionary in the reverse order of frequency and display the dictionary slice of the first 10 items.]	CO3	PO5	PSO1
5	Develop a program to read 6 subject marks from the keyboard for a student. Generate a report that displays the marks from the highest to the lowest score attained by the student. [Read the marks into a 1-Dimensional array and sort using the Bubble Sort technique].	CO1	PO2	PSO1
6	Develop a program to sort the contents of a text file and write the sorted contents into a separate text file. [Hint: Use string methods strip(), len(), list methods sort(), append(), and file methods open(), readlines(), and write()].	CO4	PO5	PSO1
7	Develop a function named DivExp which takes TWO parameters a, b, and returns a value c ( $c=a/b$ ). Write a suitable assertion for $a>0$ in the function DivExp and raise an exception for when $b=0$ . Develop a suitable program that reads two console values and calls the function DivExp.	CO3	PO2	PSO1
8	Define a function that takes TWO objects representing complex numbers and returns a new complex number with the sum of two complex numbers. Define a suitable class 'Complex' to represent the complex number. Develop a program to read N ( $N \geq 2$ ) complex numbers and to compute the addition of N complex numbers.	CO5	PO1	PSO1
9	Text Analysis Tool: Build a tool that analyses a paragraph: frequency of each word, longest word, number of sentences, etc.	CO3	PO5	PSO1
10	Develop Data Summary Generator: Read a CSV file (like COVID data or weather stats), convert to dictionary form, and allow the user to run summary queries: max, min, average by column.	CO3	PO5	PSO1
11	Develop Student Grade Tracker: Accept multiple students' names and marks. Store them in a list of tuples or dictionaries. Display summary reports (average, topper, etc.).	CO2	PO1	PSO1
12	Develop a program to display contents of a folder recursively (Directory) having sub-folders and files (name and type).	CO4	PO5	PSO1

**EXPERIMENT WISE LESSON PLAN**

S.No	Details
1a	<b>Program:</b> Develop a python program to read 2 numbers from the keyboard and perform the basic arithmetic operations based on the choice. (1-Add, 2-Subtract, 3-Multiply, 4-Divide). <b>Objective:</b> Students will be able to understand basic Python syntax, take input from the user and

	implement conditional statements to perform arithmetic operations.
1b	<b>Program:</b> Develop a program to read the name and year of birth of a person. Display whether the person is a senior citizen or not. <b>Objective:</b> Students will be able to apply input operations and conditional statements to implement logical decision making.
2a	<b>Program:</b> Develop a program to generate Fibonacci sequence of length (N). Read N from the console. <b>Objective:</b> Students will be able to understand looping constructs and generate numerical sequences using iterative programming techniques.
2b	<b>Program:</b> Write a python program to create a list and perform the following operations: Inserting an element, Removing an element, Appending an element, Displaying the length of the list, Popping an element, Clearing the list. <b>Objective:</b> Students will be able to understand Python list data structures and perform various list manipulation operations.
3a	<b>Program:</b> Read N numbers from the console and create a list. Develop a program to print mean, variance and standard deviation with suitable messages. <b>Objective:</b> Students will be able to perform statistical calculations on datasets using Python lists and arithmetic operations.
3b	<b>Program:</b> Read a multi-digit number (as chars) from the console. Develop a program to print the frequency of each digit with a suitable message. <b>Objective:</b> Students will be able to analyse numerical data and apply counting logic using loops and data structures.
4	<b>Program:</b> Develop a program to print 10 most frequently appearing words in a text file. [Hint: Use a dictionary with distinct words and their frequency of occurrences. Sort the dictionary in the reverse order of frequency and display the dictionary slice of the first 10 items.] <b>Objective:</b> Students will be able to use dictionaries and file handling techniques to analyse textual data.
5	<b>Program:</b> Develop a program to read 6 subject marks from the keyboard for a student. Generate a report that displays the marks from the highest to the lowest score attained by the student. [Read the marks into a 1-Dimensional array and sort using the Bubble Sort technique]. <b>Objective:</b> Students will be able to understand sorting algorithms and implement Bubble Sort for ordering data.
6	<b>Program:</b> Develop a program to sort the contents of a text file and write the sorted contents into a separate text file. [Hint: Use string methods strip(), len(), list methods sort(), append(), and file methods open(), readlines(), and write()]. <b>Objective:</b> Students will be able to perform file operations and manipulate textual data using Python string and list methods.
7	<b>Program:</b> Develop a function named DivExp which takes TWO parameters a, b, and returns a value c ( $c=a/b$ ). Write a suitable assertion for $a>0$ in the function DivExp and raise an exception for when $b=0$ . Develop a suitable program that reads two console values and calls the function DivExp. <b>Objective:</b> Students will be able to create functions and apply assertion and exception handling mechanisms in Python.
8	<b>Program:</b> Define a function that takes TWO objects representing complex numbers and returns a new complex number with the sum of two complex numbers. Define a suitable class 'Complex' to represent the complex number. Develop a program to read N ( $N \geq 2$ ) complex numbers and to compute the addition of N complex numbers. <b>Objective:</b> Students will be able to understand Object Oriented Programming concepts such as classes, objects and methods in Python.
9	<b>Program:</b> Text Analysis Tool: Build a tool that analyses a paragraph: frequency of each word, longest word, number of sentences, etc.. <b>Objective:</b> Students will be able to apply text processing techniques and analyse textual data using Python data structures.
10	<b>Program:</b> Develop Data Summary Generator: Read a CSV file (like COVID data or weather stats), convert to dictionary form, and allow the user to run summary queries: max, min, average by column. <b>Objective:</b> Students will be able to read structured datasets and perform basic data analysis using Python programming techniques.
11	<b>Program:</b> Develop Student Grade Tracker: Accept multiple students' names and marks. Store them in a list of tuples or dictionaries. Display summary reports (average, topper, etc.). <b>Objective:</b> Students will be able to manage structured data using lists or dictionaries and generate analytical reports.
12	<b>Program:</b> Develop a program to display contents of a folder recursively (Directory) having sub-folders and files (name and type). <b>Objective:</b> Students will be able to understand directory traversal and recursive file handling using Python modules.

## LIST OF EXPERIMENT

S.No	Experiment Name	Page No.
1a	Develop a python program to read 2 numbers from the keyboard and perform the basic arithmetic operations based on the choice. (1-Add, 2-Subtract, 3-Multiply, 4-Divide).	71
1b	Develop a program to read the name and year of birth of a person. Display whether the person is a senior citizen or not.	72
2a	Develop a program to generate Fibonacci sequence of length (N). Read N from the console.	73
2b	Write a python program to create a list and perform the following operations: Inserting an element, Removing an element, Appending an element, Displaying the length of the list, Popping an element, Clearing the list.	74
3a	Read N numbers from the console and create a list. Develop a program to print mean, variance and standard deviation with suitable messages.	75
3b	Read a multi-digit number (as chars) from the console. Develop a program to print the frequency of each digit with a suitable message.	76
4	Develop a program to print 10 most frequently appearing words in a text file. [Hint: Use a dictionary with distinct words and their frequency of occurrences. Sort the dictionary in the reverse order of frequency and display the dictionary slice of the first 10 items.]	77
5	Develop a program to read 6 subject marks from the keyboard for a student. Generate a report that displays the marks from the highest to the lowest score attained by the student. [Read the marks into a 1-Dimensional array and sort using the Bubble Sort technique].	78
6	Develop a program to sort the contents of a text file and write the sorted contents into a separate text file. [Hint: Use string methods strip(), len(), list methods sort(), append(), and file methods open(), readlines(), and write()].	79
7	Develop a function named DivExp which takes TWO parameters a, b, and returns a value c ( $c=a/b$ ). Write a suitable assertion for $a>0$ in the function DivExp and raise an exception for when $b=0$ . Develop a suitable program that reads two console values and calls the function DivExp.	80
8	Define a function that takes TWO objects representing complex numbers and returns a new complex number with the sum of two complex numbers. Define a suitable class 'Complex' to represent the complex number. Develop a program to read N ( $N \geq 2$ ) complex numbers and to compute the addition of N complex numbers.	81
9	Text Analysis Tool: Build a tool that analyses a paragraph: frequency of each word, longest word, number of sentences, etc.	82
10	Develop Data Summary Generator: Read a CSV file (like COVID data or weather stats), convert to dictionary form, and allow the user to run summary queries: max, min, average by column.	83
11	Develop Student Grade Tracker: Accept multiple students' names and marks. Store them in a list of tuples or dictionaries. Display summary reports (average, topper, etc.).	84
12	Develop a program to display contents of a folder recursively (Directory) having sub-folders and files (name and type).	85

**LAB SAFETY & USAGE INSTRUCTIONS**

1. Students must enter the computer laboratory only during scheduled lab hours and follow the instructions of the lab instructor.
2. Maintain discipline and silence inside the laboratory to ensure a productive learning environment.
3. Use only the computer system allotted and handle all equipment carefully.
4. Do not install or delete any software or modify system settings without permission.
5. Avoid using mobile phones, external storage devices, or unrelated websites during the lab session.
6. Maintain cleanliness in the laboratory and do not bring food or drinks near the computer systems.
7. Report any hardware or software issues immediately to the lab instructor or lab assistant.
8. Properly save your work, log out, and shut down the system before leaving the laboratory.

1 a. Develop a python program to read 2 numbers from the keyboard and perform the basic arithmetic operations based on the choice. (1-Add, 2-Subtract, 3-Multiply, 4-Divide).

```
# Program: Basic Arithmetic Operations

# Step 1: Read two numbers
num1 = float(input("Enter first number: "))
num2 = float(input("Enter second number: "))

# Step 2: Show choices
print("Choose operation:")
print("1 - Add")
print("2 - Subtract")
print("3 - Multiply")
print("4 - Divide")

choice = int(input("Enter your choice (1/2/3/4): "))

# Step 3: Perform operation
if choice == 1:
    print("Result =", num1 + num2)
elif choice == 2:
    print("Result =", num1 - num2)
elif choice == 3:
    print("Result =", num1 * num2)
elif choice == 4:
    if num2 != 0:
        print("Result =", num1 / num2)
    else:
        print("Error! Division by zero not allowed.")
else:
    print("Invalid choice! Please enter 1, 2, 3, or 4.")
```

```
Enter first number: 8
Enter second number: 4
Choose operation:
1 - Add
2 - Subtract
3 - Multiply
4 - Divide
Enter your choice (1/2/3/4): 1
Result = 12.0
```

```
Enter first number: 15
Enter second number: 7
Choose operation:
1 - Add
2 - Subtract
3 - Multiply
4 - Divide
Enter your choice (1/2/3/4): 2
Result = 8.0
```

1.b. Develop a program to read the name and year of birth of a person. Display whether the person is a senior citizen or not.

```
# Program: Check if person is a Senior Citizen

# Step 1: Input
name = input("Enter your name: ")
year_of_birth = int(input("Enter your year of birth: "))

# Step 2: Calculate age
current_year = 2025 # you can change this if needed
age = current_year - year_of_birth

# Step 3: Check senior citizen
if age >= 60:
    print(name, "is a Senior Citizen. Age:", age)
else:
    print(name, "is NOT a Senior Citizen. Age:", age)
```

```
Enter your name: Ramesh
Enter your year of birth: 1950
Ramesh is a Senior Citizen. Age: 75
```

```
Enter your name: Priya
Enter your year of birth: 2005
Priya is NOT a Senior Citizen. Age: 20
```

2a. Develop a program to generate Fibonacci sequence of length (N). Read N from the console.

```
N = int(input("Enter how many numbers: "))  
  
a, b = 0, 1  
  
for i in range(N):  
    print(a)  
    a, b = b, a + b
```

```
Enter how many numbers: 5  
0  
1  
1  
2  
3
```

```
Enter how many numbers: 8  
0  
1  
1  
2  
3  
5  
8  
13
```

2b. Write a python program to create a list and perform the following operations

- Inserting an element
- Removing an element
- Appending an element
- Displaying the length of the list
- Popping an element
- Clearing the list

```
# Program: List Operations

# Step 1: Create a list
my_list = [10, 20, 30]
print("Initial list:", my_list)

# Step 2: Insert an element at position 1
my_list.insert(1, 15)
print("After inserting 15 at position 1:", my_list)

# Step 3: Remove an element (value 20)
my_list.remove(20)
print("After removing 20:", my_list)

# Step 4: Append (add at end) an element
my_list.append(40)
print("After appending 40:", my_list)

# Step 5: Display length of the list
print("Length of list:", len(my_list))

# Step 6: Pop (remove last element)
my_list.pop()
print("After popping last element:", my_list)

# Step 7: Clear the list
my_list.clear()
print("After clearing list:", my_list)
```

```
Initial list: [10, 20, 30]
After inserting 15 at position 1: [10, 15, 20, 30]
After removing 20: [10, 15, 30]
After appending 40: [10, 15, 30, 40]
Length of list: 4
After popping last element: [10, 15, 30]
After clearing list: []
```

3a. Read N numbers from the console and create a list. Develop a program to print mean, variance and standard deviation with suitable messages.

**# Program: Mean, Variance, and Standard Deviation**

**# Step 1: Read N numbers**

```
N = int(input("Enter how many numbers: "))
numbers = []
```

```
for i in range(N):
    num = float(input("Enter number: "))
    numbers.append(num)
```

```
print("Numbers entered:", numbers)
```

**# Step 2: Calculate Mean**

```
mean = sum(numbers) / N
```

```
print("Mean =", mean)
```

**# Step 3: Calculate Variance**

```
variance = sum((x - mean) ** 2 for x in numbers) / N
```

```
print("Variance =", variance)
```

**# Step 4: Calculate Standard Deviation**

```
std_dev = variance ** 0.5
```

```
print("Standard Deviation =", std_dev)
```

```
Enter how many numbers: 5
Enter number: 10
Enter number: 20
Enter number: 30
Enter number: 40
Enter number: 50
Numbers entered: [10.0, 20.0, 30.0, 40.0, 50.0]
Mean = 30.0
Variance = 200.0
Standard Deviation = 14.142135623730951
```

## METHOD 2

```
import statistics
```

**# Step 1: Read N numbers**

```
N = int(input("Enter how many numbers: "))
numbers = []
```

```
for i in range(N):
    num = float(input("Enter number: "))
    numbers.append(num)
```

```
print("Numbers entered:", numbers)
```

**# Step 2: Use statistics module**

```
mean = statistics.mean(numbers)
```

```
variance = statistics.variance(numbers) # by default sample variance
```

```
std_dev = statistics.stdev(numbers) # sample standard deviation
```

```
print("Mean =", mean)
```

```
print("Variance =", variance)
```

```
print("Standard Deviation =", std_dev)
```

3b. Read a multi-digit number (as chars) from the console. Develop a program to print the frequency of each digit with a suitable message.

```
# Program: Frequency of each digit in a number

# Step 1: Read number as string (chars)
number = input("Enter a multi-digit number: ")

# Step 2: Count frequency of each digit
for digit in "0123456789": # check each digit
    count = number.count(digit)
    if count > 0:
        print("Digit", digit, "appears", count, "times")
```

```
Enter a multi-digit number: 1223455
Digit 1 appears 1 times
Digit 2 appears 2 times
Digit 3 appears 1 times
Digit 4 appears 1 times
Digit 5 appears 2 times
```

```
Enter a multi-digit number: 100202
Digit 0 appears 3 times
Digit 1 appears 1 times
Digit 2 appears 2 times
```

4. Develop a program to print 10 most frequently appearing words in a text file.  
[Hint: Use a dictionary with distinct words and their frequency of occurrences.  
Sort the dictionary in the reverse order of frequency and display the dictionary slice of the first 10 items.

```
# Program: 10 Most Frequent Words in a Text File

# Step 1: Open and read the file
filename = "sample.txt" # make sure this file exists in the same folder
with open(filename, "r") as f:
    text = f.read()

# Step 2: Split text into words
words = text.split()

# Step 3: Count frequency using dictionary
freq = {}

for word in words:
    word = word.lower() # make all words lowercase
    word = word.strip(",.!?" ) # remove punctuation
    if word in freq:
        freq[word] += 1
    else:
        freq[word] = 1

# Step 4: Sort dictionary by frequency (highest first)
sorted_freq = sorted(freq.items(), key=lambda x: x[1], reverse=True)

# Step 5: Print top 10 words
print("Top 10 most frequent words:")
for word, count in sorted_freq[:10]:
    print(word, ":", count)
```

**sample.txt contains: Python is easy to learn. Python is powerful. Python is popular and Python is fun.**

```
Top 10 most frequent words:
python : 4
is : 4
easy : 1
to : 1
learn : 1
powerful : 1
popular : 1
and : 1
fun : 1
```

5. Develop a program to read 6 subject marks from the keyboard for a student. Generate a report that displays the marks from the highest to the lowest score attained by the student. [Read the marks into a 1-Dimensional array and sort using the Bubble Sort technique].

```
# Program to sort 6 subject marks using Bubble Sort

# Read 6 marks from the user
marks = []
for i in range(6):
    mark = int(input("Enter mark: "))
    marks.append(mark)

# Bubble Sort (highest to lowest)
for i in range(len(marks)-1):
    for j in range(len(marks)-1):
        if marks[j] < marks[j+1]: # swap if smaller
            marks[j], marks[j+1] = marks[j+1], marks[j]

# Display result
print("Marks from highest to lowest:", marks)
```

```
Enter mark: 50
Enter mark: 90
Enter mark: 70
Enter mark: 85
Enter mark: 60
Enter mark: 95
Marks from highest to lowest: [95, 90, 85, 70, 60, 50]
```

6. Develop a program to sort the contents of a text file and write the sorted contents into a separate text file. [Hint: Use string methods `strip()`, `len()`, list methods `sort()`, `append()`, and file methods `open()`, `readlines()`, and `write()`].

```
# Program to sort the contents of a text file

# Step 1: Open the input file in read mode
file_in = open("input.txt", "r")

# Step 2: Read all lines from the file
lines = file_in.readlines()

# Step 3: Close the input file (good habit)
file_in.close()

# Step 4: Create an empty list to store clean lines
clean_lines = []

# Step 5: Use strip() to remove unwanted spaces/newlines
for line in lines:
    clean_line = line.strip()

    # Check length of line using len()
    if len(clean_line) > 0:      # avoid empty lines
        clean_lines.append(clean_line)  # add to list using append()

# Step 6: Sort the list of lines
clean_lines.sort()

# Step 7: Open the output file in write mode
file_out = open("output.txt", "w")

# Step 8: Write each line back to the new file
for line in clean_lines:
    file_out.write(line + "\n")

# Step 9: Close the output file
file_out.close()

print("Sorting complete! Check 'output.txt'")
```

If input.txt contains:

```
banana
apple
mango
grapes
```

Then output.txt will have:

```
apple
banana
grapes
mango
```

7. Develop a function named DivExp which takes TWO parameters a, b, and returns a value c ( $c=a/b$ ). Write a suitable assertion for  $a>0$  in the function DivExp and raise an exception for when  $b=0$ . Develop a suitable program that reads two console values and calls the function DivExp.

```
def DivExp(a, b):
    # Assertion for a > 0
    assert a > 0, "a must be greater than 0"

    # Raise error if b = 0
    if b == 0:
        raise Exception("b cannot be zero")

    # Do division
    c = a / b
    return c

# Main program
a = int(input("Enter value of a: "))
b = int(input("Enter value of b: "))

result = DivExp(a, b)
print("Result is:", result)
```

```
Enter value of a: 10
Enter value of b: 2
Result is: 5.0
```

If  $a = -5 \rightarrow$

```
AssertionError: a must be greater than 0
```

If  $b = 0 \rightarrow$

```
Exception: b cannot be zero
```

8. Define a function that takes TWO objects representing complex numbers and returns a new complex number with the sum of two complex numbers. Define a suitable class 'Complex' to represent the complex number. Develop a program to read N (N >=2) complex numbers and to compute the addition of N complex numbers.

```
# Define a class to represent Complex numbers
class Complex:
    def __init__(self, real, imag):
        self.real = real    # real part
        self.imag = imag    # imaginary part

    # Function to add two complex numbers
    def add(self, other):
        new_real = self.real + other.real
        new_imag = self.imag + other.imag
        return Complex(new_real, new_imag)

    # Function to display complex number nicely
    def show(self):
        print(f"{self.real} + {self.imag}i")

# Main Program
N = int(input("How many complex numbers? (N>=2): "))

# Read first complex number
r = int(input("Enter real part of number 1: "))
i = int(input("Enter imaginary part of number 1: "))
result = Complex(r, i)    # start with first number

# Read and add the remaining complex numbers
for k in range(2, N+1):
    r = int(input(f"Enter real part of number {k}: "))
    i = int(input(f"Enter imaginary part of number {k}: "))
    c = Complex(r, i)
    result = result.add(c)    # keep adding

# Show final result
print("Sum of all complex numbers = ", end="")
result.show()
```

```
How many complex numbers? (N>=2): 3
Enter real part of number 1: 2
Enter imaginary part of number 1: 3
Enter real part of number 2: 1
Enter imaginary part of number 2: 4
Enter real part of number 3: 3
Enter imaginary part of number 3: -2
Sum of all complex numbers = 6 + 5i
```

**9. Text Analysis Tool: Build a tool that analyses a paragraph: frequency of each word, longest word, number of sentences, etc.**

```
# Text Analysis Tool

# Step 1: Take a paragraph from user
text = input("Enter a paragraph: ")

# Step 2: Count sentences
# Sentences usually end with . or ! or ?
sentences = text.count(".") + text.count("!") + text.count("?")

# Step 3: Break the paragraph into words
words = text.split() # split() separates text at spaces

# Step 4: Count total number of words
total_words = len(words) # len() gives how many items in list

# Step 5: Find the longest word
longest = max(words, key=len) # max() finds word with biggest length

# Step 6: Show results
print("\n--- Analysis Result ---")
print("Number of sentences:", sentences)
print("Number of words:", total_words)
print("Longest word:", longest)
```

**Input:**

Enter a paragraph: Python **is** easy. Python **is** fun!

**Output**

```
--- Analysis Result ---
Number of sentences: 2
Number of words: 5
Longest word: Python
```

10. Develop Data Summary Generator: Read a CSV file (like COVID data or weather stats), convert to dictionary form, and allow the user to run summary queries: max, min, average by column.

```
import csv

# Step 1: Read CSV file into a list of dictionaries
filename = input("Enter CSV file name (example: data.csv): ")

data = []
with open(filename, "r") as f:
    reader = csv.DictReader(f)    # reads CSV as dictionary
    for row in reader:
        data.append(row)

# Step 2: Show available columns
print("Available columns:", list(data[0].keys()))

# Step 3: Ask user which column to summarize
col = input("Enter column name to summarize: ")

# Convert that column values into numbers
values = [float(row[col]) for row in data if row[col] != ""]

# Step 4: Compute summaries
maximum = max(values)
minimum = min(values)
average = sum(values) / len(values)

# Step 5: Show results
print("\n--- Summary for column:", col, "---")
print("Max:", maximum)
print("Min:", minimum)
print("Average:", average)
```

```
Day,Temp,Rainfall
1,30,5
2,32,0
3,28,12
4,35,0
```

```
Enter CSV file name (example: data.csv): weather.csv
Available columns: ['Day', 'Temp', 'Rainfall']
Enter column name to summarize: Temp
```

```
--- Summary for column: Temp ---
Max: 35.0
Min: 28.0
Average: 31.25
```

11. Develop Student Grade Tracker: Accept multiple students' names and marks. Store them in a list of tuples or dictionaries. Display summary reports (average, topper, etc.).

```
# Step 1: Read student data
n = int(input("How many students? "))

names = []
marks = []

for i in range(n):
    name = input("Enter name: ")
    mark = int(input("Enter marks: "))
    names.append(name)
    marks.append(mark)

# Step 2: Calculate average
total = 0
for m in marks:
    total = total + m
average = total / n

# Step 3: Find topper (highest marks using loop)
topper_name = names[0]
topper_marks = marks[0]

for i in range(1, n):
    if marks[i] > topper_marks:
        topper_marks = marks[i]
        topper_name = names[i]

# Step 4: Show results
print("\n--- Report ---")
print("Average marks:", average)
print("Topper:", topper_name, "with", topper_marks)
```

How many student? 3

```
Enter name: Asha
Enter marks: 80
Enter name: Ravi
Enter marks: 92
Enter name: Meena
Enter marks: 75
```

```
--- Report ---
Average marks: 82.33333333333333
Topper: Ravi with 92
```

12. Develop a program to display the contents of a folder recursively (Directory), having sub-folders and files (name and type).

```
import os

# Function to display folder contents
def show_folder(path, level=0):
    # list all items inside path
    items = os.listdir(path)
    for item in items:
        full_path = os.path.join(path, item) # complete path
        if os.path.isdir(full_path): # if it's a folder
            print("    " * level + "[Folder] " + item)
            show_folder(full_path, level + 1) # recursive call
        else: # if it's a file
            print("    " * level + "[File] " + item)

# Main Program
folder = input("Enter folder path: ")
show_folder(folder)
```

**For a folder like this:**

```
MyFolder
├── notes.txt
├── report.docx
├── Projects
│   ├── code.py
│   └── data.csv
```

**The program will print:**

```
[File] notes.txt
[File] report.docx
[Folder] Projects
    [File] code.py
    [File] data.csv
```